

REPRESENTATIONS OF COMPLEX HYPERBOLIC LATTICES INTO RANK 2 CLASSICAL LIE GROUPS OF HERMITIAN TYPE

VINCENT KOZIARZ AND JULIEN MAUBON

ABSTRACT. Let Γ be a torsion-free uniform lattice of $SU(m, 1)$, $m > 1$. Let G be either $SU(p, 2)$ with $p \geq 2$, $Sp(2, \mathbb{R})$ or $SO(p, 2)$ with $p \geq 3$. The symmetric spaces associated to these G 's are the classical bounded symmetric domains of rank 2, with the exceptions of $SO^*(8)/U(4)$ and $SO^*(10)/U(5)$. Using the correspondence between representations of fundamental groups of Kähler manifolds and Higgs bundles we study representations of the lattice Γ into G . We prove that the Toledo invariant associated to such a representation satisfies a Milnor-Wood type inequality and that in case of equality necessarily $G = SU(p, 2)$ with $p \geq 2m$ and the representation is reductive, faithful, discrete, and stabilizes a copy of complex hyperbolic space (of maximal possible induced holomorphic sectional curvature) holomorphically and totally geodesically embedded in the Hermitian symmetric space $SU(p, 2)/S(U(p) \times U(2))$, on which it acts cocompactly.

1. INTRODUCTION

Let Γ be a (torsion free) uniform lattice in the Lie group $SU(m, 1)$. We are interested here in representations, i.e. homomorphisms, of Γ in a Lie group G of Hermitian type, that is a connected semisimple Lie group with finite center and no compact factor whose associated symmetric space $\mathcal{X} = G/K$ is Hermitian (K is a maximal compact subgroup of G). We will always assume that G , hence \mathcal{X} , are irreducible. The classical groups of Hermitian type are $SU(p, q)$, $Sp(n, \mathbb{R})$, $SO^*(2n)$ and $SO_0(n, 2)$ whose associated symmetric spaces' real ranks are respectively $\min(p, q)$, n , $[n/2]$, and 2.

The Toledo invariant is a number naturally associated to such a representation $\rho : \Gamma \longrightarrow G$, and it has been recognized over the years to play a fundamental role. It is defined as follows. Let $\mathbb{H}_{\mathbb{C}}^m = SU(m, 1)/S(U(m) \times U(1))$ be complex hyperbolic m -space, and let f be any (smooth) ρ -equivariant map $\mathbb{H}_{\mathbb{C}}^m \longrightarrow \mathcal{X}$. The symmetric space \mathcal{X} being Hermitian, it is a Kähler manifold, and its G -invariant Kähler form $\omega_{\mathcal{X}}$ may be pulled-back by f to give a 2-form $f^*\omega_{\mathcal{X}}$ on $\mathbb{H}_{\mathbb{C}}^m$ which goes down by ρ -equivariance to a form on the closed complex hyperbolic manifold $M = \Gamma \backslash \mathbb{H}_{\mathbb{C}}^m$. We will make no difference between Γ -invariant objects on $\mathbb{H}_{\mathbb{C}}^m$ and the corresponding objects on M . For example, $f^*\omega_{\mathcal{X}}$ is either a 2-form on $\mathbb{H}_{\mathbb{C}}^m$ or a 2-form on M , depending on the context. Similarly, we denote by g and ω the invariant metric and Kähler form of $\mathbb{H}_{\mathbb{C}}^m$, as well as the induced metric and Kähler form on M . Now, the de Rham cohomology class in $H_{dR}^2(M)$ defined by $f^*\omega_{\mathcal{X}}$ depends only on ρ , not on f , and will be denoted by $[\rho^*\omega_{\mathcal{X}}]$. This class is then evaluated against the Kähler class of M to give the desired number

$$\tau(\rho) = \frac{1}{2m} \int_M \langle \rho^*\omega_{\mathcal{X}}, \omega \rangle dV = \frac{1}{m!} \int_M \rho^*\omega_{\mathcal{X}} \wedge \omega^{m-1}$$

where $\rho^*\omega_{\mathcal{X}}$ is any representative of $[\rho^*\omega_{\mathcal{X}}]$, $\langle \cdot, \cdot \rangle$ is the scalar product induced by g on 2-forms and $dV = \frac{1}{m!} \omega^m$ is the Riemannian volume form of $\mathbb{H}_{\mathbb{C}}^m$ (or M).

The Toledo invariant is of particular interest because

- (1) it is constant on connected components of the space of representations $\text{Hom}(\Gamma, G)$;

(2) it satisfies a Milnor-Wood type inequality, namely $\tau(\rho)$ is bounded in absolute value by a quantity depending only on the (real) rank of the symmetric space \mathcal{X} and the volume of $M = \Gamma \backslash \mathbb{H}_{\mathbb{C}}^m$. More precisely, if the Riemannian metrics on $\mathbb{H}_{\mathbb{C}}^m$ and \mathcal{X} are normalized so that the minimum of their holomorphic sectional curvatures is -1 (so that the holomorphic sectional curvature of \mathcal{X} is pinched between -1 and $-1/\text{rk } \mathcal{X}$), the following holds:

$$|\tau(\rho)| \leq \text{rk } \mathcal{X} \text{Vol}(M).$$

(3) maximal representations, i.e. representations ρ for which $|\tau(\rho)| = \text{rk } \mathcal{X} \text{Vol}(M)$, are expected to be of a very special kind, and therefore rigidity results should follow.

The Toledo invariant appeared for the first time in 1979 Toledo's paper [To79] and more explicitly in [To89], where the Milnor-Wood inequality (2) was proved for $m = 1$ and $\text{rk } \mathcal{X} = 1$, namely Γ is the fundamental group of a Riemann surface and $G = \text{SU}(n, 1)$. Toledo also proved that maximal representations are faithful with discrete image, and stabilize a complex line in the complex hyperbolic n -space \mathcal{X} , generalizing Goldman's results for $G = \text{SL}(2, \mathbb{R})$ [Go80, Go88]. At approximately the same time, Corlette, using a very similar invariant, the volume of the representation, established in [Co88] the same kind of result for $m \geq 2$ and $G = \text{SU}(n, 1)$. An immediate corollary is that a uniform lattice in $\text{SU}(m, 1)$ can not be deformed non-trivially in $\text{SU}(n, 1)$, $n \geq m \geq 1$, a result first obtained by Goldman and Millson in [GM87] using different methods. These results have been extended to the non-uniform case in [BI07, KM08] (the definition of the Toledo invariant must be modified). Therefore the case where the rank of the symmetric space associated to G is 1 is now settled.

Using the work of Domic-Toledo [DT87] and Clerc-Ørsted [CØ01], Burger and Iozzi obtained in [BI07] the Milnor-Wood inequality (2) in full generality. Since then, much progress has been made and maximal representations of the fundamental group of a Riemann surface into such general groups of Hermitian type are well understood [BIW03, BIW06, BGG03, BGG05].

So far the expected rigidity for maximal representations of higher dimensional complex hyperbolic lattices in Lie groups of Hermitian type of rank at least two has not been proved and this is the question we (begin to) address in this paper. Since the Milnor-Wood inequality (2) is known, one would a priori like to focus on point (3) and give a complete description of the maximal representations. Our strategy will be different, and to explain why we need to say a word about the available methods to prove the results we mentioned. Essentially, there are two different ways of attacking the problem. The one used by Burger and Iozzi relies on bounded cohomology theory and allows to prove the bound (2) in great generality but gives relatively few informations on the maximal representations, so that a separate study has to be made. The second one, used in [Co88, KM08] and in [BGG03, BGG05], relies on harmonic maps and/or Higgs bundles machinery and belongs more to the world of complex differential geometry. Following this approach, as we shall, makes it quite difficult to prove the Milnor-Wood inequality (and in fact no such proof is known in the general case) but once it is proved (in some special cases), maximal representations are easier to understand.

The Higgs bundle theory was developed by Hitchin [Hi87, Hi92] for Riemann surfaces and Simpson [Si88, Si92, Si94a, Si94b] in higher dimensions to study fundamental groups of Kähler manifolds and their linear representations. It consists in establishing a correspondence between representations of the fundamental group Γ of a Kähler manifold M and purely holomorphic objects, called polystable Higgs bundles over M , and then working on these objects with tools from complex geometry. This is in fact only possible for reductive representations, since the construction of Higgs bundles requires harmonic maps, but this restriction will not be a serious issue for our purposes. We will explain this correspondence in some details in section 2. For now let us simply say that to a reductive representation of Γ is associated a Higgs bundle on

M , that is a holomorphic vector bundle $(E, \bar{\partial})$ on M together with a holomorphic $(1,0)$ -form θ , the Higgs field, which takes values in (a subbundle of) the bundle of endomorphisms of E and satisfies $[\theta, \theta] = 0$. We shall often write $\theta : E \rightarrow E \otimes \Omega^1$, where Ω^1 is the sheaf (of germs) of holomorphic 1-forms on M , which we identify with the holomorphic cotangent bundle of M . Polystability is a condition that relates the slope of proper θ -invariant saturated subsheaves of E to the slope of E itself (which will be 0 here). Recall that the slope of a saturated sheaf is defined as the quotient of its degree by its rank.

The Toledo invariant may be defined as before for representations ρ of the fundamental group Γ of any closed Kähler manifold M in a Lie group of Hermitian type G (f is then simply a ρ -equivariant map from the universal cover \widetilde{M} of M to $\mathcal{X} = G/K$), but in the Higgs bundles setting it is best interpreted as the degree of a complex vector bundle over M as follows. The Hermitian symmetric space \mathcal{X} is Kähler-Einstein and therefore $\omega_{\mathcal{X}}$ is up to a constant the first Chern class of the holomorphic tangent bundle $T\mathcal{X}$ of \mathcal{X} . Therefore $f^*\omega_{\mathcal{X}}$ is up to a constant the first Chern class of the induced bundle $f^*T\mathcal{X}$ over M so that $\tau(\rho)$ is, again up to a constant, simply the degree of this bundle. To be more precise, if the Ricci curvature tensor of \mathcal{X} is $\lambda_{\mathcal{X}} g_{\mathcal{X}}$ (remember that the Riemannian metric of \mathcal{X} is normalized to have minimal holomorphic sectional curvature -1), we have $\tau(\rho) = -\frac{2\pi}{m!\lambda_{\mathcal{X}}} \deg f^*T\mathcal{X}$. We remark that if M is complex hyperbolic of complex dimension m , $\text{Vol}(M) = \frac{4\pi}{m!(m+1)} \deg \Omega^1$. Hence the Milnor-Wood type bound (2) can be written:

$$(2') \quad \left| \frac{\deg f^*T\mathcal{X}}{2\lambda_{\mathcal{X}}} \right| \leq \text{rk } \mathcal{X} \frac{\deg \Omega^1}{m+1},$$

an interpretation we will use constantly. If the ρ -equivariant map $f : \widetilde{M} \rightarrow \mathcal{X}$ is chosen to be harmonic, the bundle $f^*T\mathcal{X} \rightarrow M$ is constructed from holomorphic subbundles of the Higgs bundle $E \rightarrow M$ associated to the representation ρ . Moreover, the fact that ρ is not valued in the full general linear group but in a smaller group G of Hermitian type implies that the Higgs bundle E has a special structure. The idea is then that Inequality (2') will follow from this particular structure and the polystability condition. This implementation of the Higgs bundle theory has been carried out for Riemann surfaces, for example by Xia [Xi00] in some special cases, and more generally by Bradlow, Garcia-Prada and Gothen in [BGG03, BGG05]. The Kähler manifold M being a complex curve in their situation makes it quite easy to deduce (2') from the structure of the Higgs bundles and maximal representations can be studied in great details. It is for example possible to count the number of connected components of the moduli space of maximal representations. The reader should consult the papers [BGG03, BGG05] to see the strength of the method in this case.

When $M = \Gamma \backslash \mathbb{H}_{\mathbb{C}}^m$ is a (closed) complex hyperbolic manifold of dimension $m \geq 2$, one expects maximal representations to be extremely restricted. In fact they should all be induced by special holomorphic or antiholomorphic totally geodesic embeddings of complex hyperbolic m -space into the Hermitian symmetric space \mathcal{X} associated to G .

Inequality (2) or (2') for such higher dimensional lattices turns out to be surprisingly difficult to prove. In particular, it seems necessary to first deform the Higgs bundle associated to the representation to a more tractable one. This is done via the \mathbb{C}^* -action on the moduli space of Higgs bundles, see section 2.4 for details. In the general case, the structure of the systems of Hodge bundles obtained in this way is still very complicated. However, when the rank of the symmetric space \mathcal{X} is 2, they take simpler forms which, together with the crucial fact that the holomorphic tangent bundle of a complex hyperbolic manifold is stable (since such a manifold is Kähler-Einstein), allow us to prove our main result:

Theorem 1.1. *Let Γ be a (torsion-free) uniform lattice in $SU(m, 1)$. Let G be either $SU(p, 2)$ with $p \geq 2$, $SO_0(p, 2)$ with $p \geq 3$ or $Sp(2, \mathbb{R})$. Finally let $\rho : \Gamma \rightarrow G$ be a representation.*

Then $|\tau(\rho)| \leq 2\text{Vol}(\Gamma \backslash \mathbb{H}_{\mathbb{C}}^m)$. If $m > 1$ and ρ is maximal, namely if $|\tau(\rho)| = 2\text{Vol}(\Gamma \backslash \mathbb{H}_{\mathbb{C}}^m)$, then $G = SU(p, 2)$ with $p \geq 2m$, ρ is reductive, faithful, discrete, and stabilizes a holomorphic totally geodesic copy of complex hyperbolic m -space of holomorphic sectional curvature $-1/2$ in the Hermitian symmetric space $\mathcal{X} = SU(p, 2)/S(U(p) \times U(2))$. Moreover Γ acts cocompactly on this copy of complex hyperbolic space.

Although Higgs bundles are associated to reductive representations, we do not assume that ρ is reductive in the theorem. This is because every representation can be deformed to a reductive one, an operation that does not change the value of the Toledo invariant. Moreover we shall see in section 3.3.3 that non reductive representations can not be maximal.

The theorem covers all but two of the classical Lie groups of Hermitian type G whose associated symmetric spaces' rank is 2. The missing ones are $SO^*(8)$ and $SO^*(10)$. In fact, $SO^*(8)$ and $SO_0(6, 2)$ are isogenous and have the same associated symmetric space. This means that if a representation $\Gamma \rightarrow SO^*(8)$ lifts to $Spin(6, 2)$, then projecting down to $SO_0(6, 2)$ gives the result for this representation as well.

The proof of the Milnor-Wood type inequality $|\tau(\rho)| \leq 2\text{Vol}(M)$ is given in Section 3 for $G = SU(p, 2)$ and in Section 4 for $G = SO_0(p, 2)$. The case of $Sp(2, \mathbb{R})$ follows from the case of $SU(2, 2)$ since $Sp(2, \mathbb{R}) \subset SU(2, 2)$.

The theorem in particular says that for $m > 1$ there is no maximal representations of a uniform lattice Γ of $SU(m, 1)$ in $SO_0(p, 2)$, $Sp(2, \mathbb{R})$ or $SU(p, 2)$ with $p < 2m$. Our method indeed yields explicit better bounds on the Toledo invariant in these cases. For representations $\rho : \Gamma \rightarrow SU(p, 2)$, the arguments of [VZ07] can be adapted to give the following, which is stronger than the Milnor-Wood inequality (2) exactly when $p < 2m$:

Proposition 1.2. *Let Γ be a (torsion-free) uniform lattice in $SU(m, 1)$ and let $\rho : \Gamma \rightarrow SU(p, 2)$, $p \geq 2$, be a representation. Then $|\tau(\rho)| \leq \frac{2p}{p+2} \frac{m+1}{m} \text{Vol}(\Gamma \backslash \mathbb{H}_{\mathbb{C}}^m)$.*

If $\rho : \Gamma \rightarrow G$ is a maximal representation, for which as we said $G = SU(p, 2)$ with $p \geq 2m$, we will prove that there exists a maximal holomorphic or antiholomorphic totally geodesic ρ -equivariant embedding $\mathbb{H}_{\mathbb{C}}^m \rightarrow \mathcal{X} = SU(p, 2)/S(U(p) \times U(2))$, from which the assertions of our main theorem follow. By a maximal embedding $\mathbb{H}_{\mathbb{C}}^m \rightarrow \mathcal{X}$ we mean an embedding whose image's induced holomorphic sectional curvature is everywhere the greatest possible, namely $-1/2$ with our normalization. See section 3.1.2 for a discussion of these embeddings and a description of the stabilizer in $SU(p, 2)$ of their images in \mathcal{X} . If $f : \mathbb{H}_{\mathbb{C}}^m \rightarrow \mathcal{X}$ is such a ρ -equivariant maximal embedding, we will loosely say that ρ is induced by f , although f determines $\rho(\gamma)$ for $\gamma \in \Gamma$ only up to composition with an element of $SU(p, 2)$ fixing pointwise the image of f in \mathcal{X} .

The paper is organized as follows. Section 2 is an overview on how Higgs bundles are constructed from representations of the fundamental group of a Kähler manifold. We say a few words about the corresponding moduli space, the \mathbb{C}^* -action it comes with, and the systems of Hodge bundles that are obtained as fixed points of this action. Section 3 is devoted to the proof of the main theorem when the representation takes values in $SU(p, 2)$, which is the most interesting case. The first subsection is expository, we give there the necessary background on the geometry of the associated Hermitian symmetric space. This is used in the next subsection to describe the particular structure of the Higgs bundles associated to such a representation. The third subsection contains the proof of the Milnor-Wood type inequality and the fourth deals with maximal representations. In the fifth we prove Proposition 1.2. Finally, Section 4 follows the lines of Section 3 in the case of $SO_0(p, 2)$: the first subsection

describes the associated symmetric space whereas the second is devoted to the Higgs bundles arising in this case and to the proof of the Milnor-Wood type inequality.

Acknowledgments. We would like to thank Jean-Louis Clerc and Andrei Teleman for useful discussions and their interest in our work.

2. REPRESENTATIONS OF THE FUNDAMENTAL GROUP, FLAT BUNDLES AND HIGGS BUNDLES

In this section we give a short presentation of the links between representations of the fundamental group Γ of a polarized projective variety M and Higgs bundles on M . To be a little more precise, we will explain in some details how G -Higgs bundles are constructed from reductive representations of Γ into a group G of Hodge type. There is in fact a much deeper correspondence between reductive representations of Γ in G and G -Higgs bundles over M with some stability properties. However, we shall not need the full strength of this correspondence, and we will stick to what matters for our purposes. We refer to the original papers of Simpson [Si88, Si92, Si94a, Si94b] and also to [BGG03, BGG05]. These results will be applied to the case where M is a compact complex hyperbolic manifold, with the polarization given by its Kähler form, and G is a Lie group of Hermitian type, that is, a group of Hodge type whose associated symmetric space \mathcal{X} is Hermitian.

Let M be a polarized projective variety, Γ its fundamental group, and \widetilde{M} its universal cover, so that $M = \Gamma \backslash \widetilde{M}$. Let \mathbb{E} be a finite dimensional complex vector space and G be a connected real algebraic non compact simple subgroup of $\mathrm{GL}(\mathbb{E})$. We assume that the action of G on \mathbb{E} is irreducible and that G is of Hodge type, meaning that G admits a Cartan involution c which is an inner automorphism given by an element of the connected component of the complexification $G_{\mathbb{C}}$ of G . Recall that if σ denotes complex conjugation in $G_{\mathbb{C}}$ relative to G , a Cartan involution is an involution c of G such that $\tau := c\sigma = \sigma c$ is complex conjugation with respect to a compact real form of $G_{\mathbb{C}}$ (see [Si92, pp. 46-50] for details). Let K be a maximal compact subgroup of G , so that $\mathcal{X} = G/K$ is the symmetric space associated to G . Finally, let ρ be a representation $\Gamma \rightarrow G$.

2.1. Real Higgs equations.

Let P_G be the flat principal G -bundle $\widetilde{M} \times_{\rho} G$ on M associated to the representation ρ . A metric on P_G is a reduction of the structure group G of P_G to its maximal compact subgroup K , namely, a K -principal subbundle P_K of P_G . This is the same thing as a section of the associated bundle $P_G \times_G \mathcal{X} \simeq P_G/K$ over M . In our setting, since P_G is flat, this associated bundle is isomorphic to $\widetilde{M} \times_{\rho} \mathcal{X}$, and a section of this bundle is given by a ρ -equivariant map $f : \widetilde{M} \rightarrow \mathcal{X}$. In this case, the K -principal bundle $G \rightarrow \mathcal{X} = G/K$ can be pulled-back by f to give a K -principal bundle $f^*G \subset \widetilde{M} \times G$ over \widetilde{M} . This bundle goes down under the action of Γ and yields the K -principal bundle $P_K \subset P_G$ over M . Note that P_G is recovered as the bundle $P_K \times_K G$ associated to P_K via the action of K on G by left translations.

$$\begin{array}{ccccc}
 & & f^*G \subset \widetilde{M} \times G & & G \\
 & \swarrow & \downarrow & & \downarrow \\
 P_K \subset P_G & & \widetilde{M} & \xrightarrow{f} & \mathcal{X} \\
 \downarrow & \swarrow & & & \\
 M & & & &
 \end{array}$$

Let $\tilde{\omega}_G$ be the flat connection 1-form on the trivial G -bundle $\tilde{M} \times G \rightarrow \tilde{M}$: if $X \in T\tilde{M}$ and A^* is the left invariant vector field on G corresponding to $A \in \mathfrak{g}$, $\tilde{\omega}_G(X, A^*) = A$. This form goes down under the Γ -action to give the flat connection ω_G on P_G . On the bundle $G \rightarrow \mathcal{X}$ we have the usual invariant connection λ defined by $\lambda(A^*) = A_{\mathfrak{k}}$, where $A_{\mathfrak{k}}$ is the \mathfrak{k} -component of $A \in \mathfrak{g}$ in the Cartan decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$. Let $\tilde{\omega}_K = f^*\lambda$ be the induced connection 1-form on the pull-back $f^*G \rightarrow \tilde{M}$. Again, $\tilde{\omega}_K$ is Γ -invariant and gives a connection 1-form ω_K on $P_K \rightarrow M$.

For $X \in T_x\tilde{M}$, we have $d_x f(X) \in T_{f(x)}\mathcal{X} = T_{gK}G/K = g_*T_{eK}G/K = g_*\mathfrak{p}$. Hence we can define a form $\tilde{\Theta} \in \Omega^1(\tilde{M} \times G) \otimes \mathfrak{p}$ by $\tilde{\Theta}(X, A^*) = g_*^{-1}df(X)$. If we restrict it to f^*G , we have that $\tilde{\Theta}(X, A^*) = A_{\mathfrak{p}}$. Hence, on f^*G , we have $\tilde{\omega}_G = \tilde{\omega}_K + \tilde{\Theta}$. $\tilde{\Theta}$ is Γ -invariant and gives an element Θ of $\Omega^1(P_K) \otimes \mathfrak{p}$ so that $\omega_G = \omega_K + \Theta$ on P_K . The form Θ behaves well under the right action of K on P_K : $R_k^*\Theta = \text{Ad}(k^{-1})\Theta$. Moreover Θ vanishes on vectors tangent to the fibers of $P_K \rightarrow M$. Hence Θ can be seen as a 1-form on M with values in the vector bundle $P_K \times_{\text{Ad}} \mathfrak{p} \rightarrow M$ associated to P_K via the adjoint action of K on \mathfrak{p} . One should remark that this vector bundle is nothing but the quotient under Γ of the pull-back $f^*T\mathcal{X}$ of the tangent bundle $T\mathcal{X}$ of \mathcal{X} .

Let d_G , F_G and d_K , F_K be the covariant exterior derivatives and the curvature forms of the connection 1-forms ω_G and ω_K . We have for example $d_G = d + \text{ad}(\omega_G)$ and $F_G = d\omega_G + \frac{1}{2}[\omega_G, \omega_G]$.

Since $\omega_G = \omega_K + \Theta$ is flat, we have $0 = F_G = F_K + \frac{1}{2}[\Theta, \Theta] + d_K\Theta$. Decomposing according to $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$, we obtain

$$\begin{cases} F_K + \frac{1}{2}[\Theta, \Theta] = 0 \\ d_K\Theta = 0 \end{cases}$$

Moreover, Corlette proved in [Co88] that if the representation ρ is reductive, that is if the Zariski closure of $\rho(\Gamma)$ is a reductive subgroup of G , then there exists a harmonic ρ -equivariant map $f : \tilde{M} \rightarrow \mathcal{X}$. In our setting, this means that Θ can be chosen to satisfy the additional condition

$$d_K^*\Theta = 0.$$

Summing up, we see that a reductive representation $\rho : \Gamma \rightarrow G$ is equivalent to the data of a connection d_K on a K -principal bundle $P_K \rightarrow M$ and an element $\Theta \in \Omega^1(M, P_K \times_{\text{Ad}} \mathfrak{p})$ satisfying

$$\begin{cases} F_K + \frac{1}{2}[\Theta, \Theta] = 0 \\ d_K\Theta = 0 \\ d_K^*\Theta = 0 \end{cases} \quad (\text{real Higgs equations})$$

So far all that was needed was M to be a compact manifold and $\mathcal{X} = G/K$ to be a symmetric space of the non-compact type.

2.2. Complex Higgs equations.

Let $G_{\mathbb{C}}$ be the complexification of G and $K_{\mathbb{C}}$ be the subgroup of fixed points of the Cartan involution c in $G_{\mathbb{C}}$. $K_{\mathbb{C}}$ is the complexification of K . Let also $\mathfrak{g}_{\mathbb{C}} = \mathfrak{k}_{\mathbb{C}} \oplus \mathfrak{p}_{\mathbb{C}}$ be the corresponding Cartan decomposition of the Lie algebra of $G_{\mathbb{C}}$.

The K -principal bundle P_K can be extended to a $K_{\mathbb{C}}$ -principal bundle $P_{K_{\mathbb{C}}} = P_K \times_K K_{\mathbb{C}}$. We still denote by d_K the extension of the covariant exterior derivative to $P_{K_{\mathbb{C}}}$. Θ can be extended by \mathbb{C} -linearity to an element of $\Omega_{\mathbb{C}}^1(M, P_{K_{\mathbb{C}}} \times_{\text{Ad}} \mathfrak{p}_{\mathbb{C}}) = \Omega^0(T_{\mathbb{C}}^*M \otimes (P_{K_{\mathbb{C}}} \times_{\text{Ad}} \mathfrak{p}_{\mathbb{C}}))$. Decomposing according to types we write $d_K = d_K^{1,0} + d_K^{0,1}$ and $\Theta = \Theta^{1,0} + \Theta^{0,1}$. We set $\theta = \Theta^{1,0} \in \Omega^{1,0}(M, P_{K_{\mathbb{C}}} \times_{\text{Ad}} \mathfrak{p}_{\mathbb{C}})$. We have $\Theta = \theta - \tau(\theta)$, for θ comes from a real form.

It has been shown by Sampson [Sam86] that the harmonicity of the map f and the facts that M is Kähler and \mathcal{X} has non positive complexified sectional curvature implies that f is pluriharmonic, which can be written $d_K^{0,1}\theta = 0$. Moreover the complexified sectional curvature of \mathcal{X} vanishes on the image of the $(1,0)$ -part of f , i.e. on the image of θ . Since here \mathcal{X} is a symmetric space, the curvature is given by the Lie bracket and the vanishing of the complexified sectional curvature just means that $[\theta, \theta] = 0$. From this it follows that $(d_K^{0,1})^2 = 0$, namely that $d_K^{0,1}$ defines a holomorphic structure on the principal bundle $P_{K_{\mathbb{C}}} \rightarrow M$ and on the associated complex vector bundle $P_{K_{\mathbb{C}}} \times_{\text{Ad}} \mathfrak{p}_{\mathbb{C}}$. The pluriharmonicity of f therefore means that θ is holomorphic for this holomorphic structure.

Looking at the real Higgs equations, we also obtain that $F_K - [\theta, \tau(\theta)] = 0$.

Therefore, starting from a reductive representation ρ , we have constructed a $K_{\mathbb{C}}$ -principal bundle $P_{K_{\mathbb{C}}} = P_K \times_K K_{\mathbb{C}}$ over M equipped with a complex structure $d_K^{0,1}$, and a holomorphic section θ of $\Omega^1 \otimes (P_{K_{\mathbb{C}}} \times_{\text{Ad}} \mathfrak{p}_{\mathbb{C}})$, where Ω^1 is the sheaf of holomorphic 1-forms on M , satisfying

$$\begin{cases} [\theta, \theta] = 0 \\ F_K - [\theta, \tau(\theta)] = 0 \end{cases} \quad (\text{complex Higgs equations})$$

2.3. Higgs bundles and stability conditions.

A holomorphic $K_{\mathbb{C}}$ -principal bundle $P_{K_{\mathbb{C}}}$ together with a holomorphic section θ of $\Omega^1 \otimes (P_{K_{\mathbb{C}}} \times_{\text{Ad}} \mathfrak{p}_{\mathbb{C}})$ satisfying $[\theta, \theta] = 0$ will be called a G -Higgs principal bundle on M . This is a purely holomorphic object. The section θ is called the Higgs field.

The groups $G_{\mathbb{C}}$ and $K_{\mathbb{C}}$ act on the vector space \mathbb{E} , and we can form the associated vector bundle $E = P_{K_{\mathbb{C}}} \times_{K_{\mathbb{C}}} \mathbb{E}$. The holomorphic structure on $P_{K_{\mathbb{C}}}$ induces a holomorphic structure on E . Moreover, $\mathfrak{p}_{\mathbb{C}}$ is a subspace of $\text{End}(\mathbb{E})$, hence we may consider θ as a holomorphic $(1,0)$ -form on M with values in the bundle $\text{End}(E)$ (we will write $\theta : E \rightarrow E \otimes \Omega^1$). The pair (E, θ) is called a G -Higgs (vector) bundle. If we forget the group G and consider only the data of a holomorphic vector bundle E together with a holomorphic $(1,0)$ -form θ with values in $\text{End}(E)$, we obtain what is called a Higgs (vector) bundle. A saturated subsheaf of a Higgs bundle (E, θ) is said to be a Higgs subsheaf, or a θ -invariant subsheaf, if $\theta(\mathcal{F}) \subset \mathcal{F} \otimes \Omega^1$.

Using the given polarization on M , we can compute the degree $\text{deg } \mathcal{F}$ and the slope $\mu(\mathcal{F}) = \frac{\text{deg } \mathcal{F}}{\text{rk } \mathcal{F}}$ of any saturated subsheaf \mathcal{F} of E .

A Higgs bundle (E, θ) is called stable (resp. semistable) if for every Higgs subsheaf \mathcal{F} of E such $0 < \text{rk } \mathcal{F} < \text{rk } E$, the slope $\mu(\mathcal{F})$ of \mathcal{F} is smaller (resp. not bigger) than the slope $\mu(E)$ of E . It is called polystable if it is the sum of stable Higgs bundles of the same slope. If the Higgs bundle (E, θ) is polystable and \mathcal{F} is a proper Higgs subsheaf of E with $\mu(\mathcal{F}) = \mu(E)$ then \mathcal{F} is in fact a Higgs subbundle of E and E splits as the direct sum of \mathcal{F} with another Higgs subbundle of the same slope [Si88].

If a G -Higgs principal bundle $(P_{K_{\mathbb{C}}}, \theta)$ on M comes from a reductive representation of Γ in G as above, the connection d_K on the reduction P_K of the structure group of $P_{K_{\mathbb{C}}}$ to K induces a connection on the associated G -Higgs vector bundle (E, θ) compatible with its complex structure. The curvature of this connection can be used to compute the degree (and the slope) of saturated subsheaves of E and the condition $F_K - [\theta, \tau(\theta)] = 0$ then implies that (E, θ) is polystable [Si88]. Remark also that in this case, E is flat as a C^∞ -bundle, hence its slope is zero.

In the terminology of [Si92, part 6], G -Higgs principal bundles constructed from reductive representations of Γ in G are reductive $(G_{\mathbb{C}}, c)$ -torsors. Remark also that everything until now is valid for M a compact Kähler manifold, with the polarization given by its Kähler form.

2.4. Moduli space, \mathbb{C}^* -action and systems of Hodge bundles.

Let H be a complex reductive group and M a projective variety. A H -Higgs principal bundle on M consists of a holomorphic H -principal bundle P_H over M , together with a holomorphic section θ of $\Omega^1 \otimes (P_H \times_{\text{Ad}} \mathfrak{h})$ satisfying $[\theta, \theta] = 0$. The H -Higgs principal bundle is called (semi-, poly-)stable, if for a (hence any) faithful representation \mathbb{E} of H , the corresponding H -Higgs vector bundle $E := P_H \times_H \mathbb{E}$ is (semi-, poly-)stable.

In [Si94a, Si94b], Simpson constructed the moduli space \mathcal{M}_H of equivalence classes of semistable H -Higgs principal bundles over a projective variety M . A very important feature of this moduli space is that it comes with a natural analytic \mathbb{C}^* -action given by $t.[P_H, \theta] = [P_H, t\theta]$ for $t \in \mathbb{C}^*$. Using the fact that the moduli space admits a proper map to a vector space, Simpson proved in [Si94b] that for any $[P_H, \theta] \in \mathcal{M}_H$, the limit of $[P_H, t\theta]$ as $t \in \mathbb{C}^*$ goes to zero exists and is unique. The limit is therefore a fixed point of the \mathbb{C}^* -action. A Higgs vector bundle E associated to this limit hence has the structure of a so-called system of Hodge bundles [Si88, Si92]. This means that E splits holomorphically as a sum $E_1 \oplus \dots \oplus E_k$ of holomorphic vector bundles and that the Higgs field in $\Omega^1 \otimes (P_H \times_{\text{Ad}} \mathfrak{h}) \subset \Omega^1 \otimes \text{End}E$ is given by a collection of holomorphic maps $\theta_i : E_i \rightarrow E_{i+1} \otimes \Omega^1$, $1 \leq i \leq k$ (with the convention that $E_{k+1} = \{0\}$). We will abuse notation and use the following kind of diagram:

$$E_1 \xrightarrow{\theta_1} E_2 \xrightarrow{\theta_2} \dots \xrightarrow{\theta_{k-1}} E_k \xrightarrow{\theta_k} 0$$

to denote such a system of Hodge bundles.

Now assume that we have a reductive representation of the fundamental group of M in a group of Hodge type $G < \text{GL}(\mathbb{E})$ as above, and consider the G -Higgs principal bundle $(P_{K_{\mathbb{C}}}, \theta)$ constructed in the previous paragraphs. Let also $E = P_{K_{\mathbb{C}}} \times_{K_{\mathbb{C}}} \mathbb{E}$ be the associated polystable G -Higgs vector bundle on M . The G -Higgs principal bundle $(P_{K_{\mathbb{C}}}, \theta)$ is a fortiori a semistable H -Higgs principal bundle, for $H = G_{\mathbb{C}}$. Therefore, we may deform this bundle to a fixed point of the \mathbb{C}^* -action in the moduli space $\mathcal{M}_{G_{\mathbb{C}}}$. It will be important for our purpose to know that the resulting structure of system of Hodge bundles on E is compatible with the initial structure of G -Higgs vector bundle. In order to explain what we mean by this, we formulate the conditions needed for a $G_{\mathbb{C}}$ -Higgs vector bundle E to be a G -Higgs vector bundle with the help of the Cartan involution c of $G_{\mathbb{C}}$.

Call α the element of $G_{\mathbb{C}} < \text{GL}(\mathbb{E})$ such that $c(g) = \alpha g \alpha^{-1}$, $\forall g \in G_{\mathbb{C}}$. Of course α^2 is in the center of $G_{\mathbb{C}}$. Since G is simple and is not a complex group seen as a real group (because it is of Hodge type), $G_{\mathbb{C}}$ is simple too, so that its center is finite and hence conjugated to some subgroup of a unitary group in $\text{GL}(\mathbb{E})$. Therefore α^2 is diagonalizable. The action of $G_{\mathbb{C}}$ on \mathbb{E} is irreducible, thus α^2 acts as a scalar on \mathbb{E} . We deduce that α is diagonalizable with two eigenvalues ξ and $-\xi$. There is an automorphism of E coming from α , because the structure group of E is $K_{\mathbb{C}}$ which is precisely the subgroup of elements of $G_{\mathbb{C}}$ fixed by c . Call this automorphism also α . If \mathbb{V} and \mathbb{W} are the eigenspaces corresponding to the two eigenvalues $\pm\xi$ of $\alpha \in \text{GL}(\mathbb{E})$, the bundle E splits holomorphically as the sum of the two eigenbundles $V = P_{K_{\mathbb{C}}} \times_{K_{\mathbb{C}}} \mathbb{V}$ and $W = P_{K_{\mathbb{C}}} \times_{K_{\mathbb{C}}} \mathbb{W}$ of $\alpha \in \text{Aut}(E)$. Moreover the Higgs field θ is a holomorphic section of $(\text{Hom}(V, W) \oplus \text{Hom}(W, V)) \otimes \Omega^1$ since it satisfies $\alpha \theta \alpha^{-1} = -\theta$, for it takes values in $\mathfrak{p}_{\mathbb{C}}$, which is the (-1) -eigenspace of c acting on $\mathfrak{g}_{\mathbb{C}}$.

If now (E_0, θ_0) is the polystable $G_{\mathbb{C}}$ -Higgs vector bundle obtained after deformation of (E, θ) by the \mathbb{C}^* -action, then the decomposition $V \oplus W$ is still holomorphic for the complex structure of E_0 , and $\alpha \theta_0 \alpha^{-1} = -\theta_0$ holds in the limit. Indeed, let us denote by $\bar{\partial}_V$ and $\bar{\partial}_W$ the restrictions to V and W of the complex structure $\bar{\partial}_E$ of E , and let $t_i \in \mathbb{C}^*$ be a sequence converging to 0. Since $\alpha \theta \alpha^{-1} = -\theta$, Theorem 1 in [Si88] implies that there exist harmonic metrics ([Si92]) on the bundles $(E, \bar{\partial}_V \oplus \bar{\partial}_W, t_i \theta)$ for which the decomposition $V \oplus W$

is orthogonal. From this it follows as in the proof of Lemma 2.8 in [Si92] that there are C^∞ -automorphisms φ_i and ψ_i of V and W respectively such that (subsequences of) the sequences $\varphi_i^* \bar{\partial}_V$ and $\psi_i^* \bar{\partial}_W$ converge to complex structures $\bar{\partial}_{V_0}$ and $\bar{\partial}_{W_0}$ on V and W . Moreover, if ϕ_i is the automorphism $\varphi_i \oplus \psi_i$, then (a subsequence of) the sequence $\phi_i^{-1} \circ (t_i \theta) \circ \phi_i$ converges to a Higgs field θ_0 holomorphic for the new complex structure $\bar{\partial}_{E_0} = \bar{\partial}_{V_0} \oplus \bar{\partial}_{W_0}$ on E . This is what we want since the Higgs field θ_0 then clearly verifies $\alpha \theta_0 \alpha^{-1} = -\theta_0$.

The structure of system of Hodge bundles $E_0 = E_1 \oplus \dots \oplus E_k$ is also compatible with the decomposition $V \oplus W$ in the sense that for each i , $E_i = V_i \oplus W_i$, where V_i and W_i are subbundles of V and W respectively. To see this, remember that the E_i 's are in fact the characteristic bundles of a holomorphic section ϕ of $\text{End} E_0$ such that $[\phi, \theta_0] = \sqrt{-1} \theta_0$. This follows from the fact that (E_0, θ_0) is fixed under the action of \mathbb{C}^* in the moduli space. Now, because $\alpha \theta_0 = -\theta_0 \alpha$, we may in fact choose a ϕ which commutes with α , in which case its characteristic bundles will split as desired along V and W .

To summarize, we have

Proposition 2.1. *A polystable G -Higgs vector bundle (E, θ) admits a holomorphic splitting $E = V \oplus W$ with respect to which θ is off-diagonal: θ is a holomorphic $(1,0)$ -form taking values in the bundle $\text{Hom}(V, W) \oplus \text{Hom}(W, V)$. Such a bundle can be deformed via the \mathbb{C}^* -action to a polystable system of Hodge bundles $E_1 \oplus \dots \oplus E_k$ whose Higgs field is given by holomorphic maps $\theta_i : E_i \rightarrow E_{i+1} \otimes \Omega^1$ with the following properties: for all i , there are holomorphic subbundles V_i and W_i of V and W respectively, such that $E_i = V_i \oplus W_i$, and θ_i maps V_i to $W_{i+1} \otimes \Omega^1$ and W_i to $V_{i+1} \otimes \Omega^1$.*

Remark. It would be more satisfactory to deduce all this from the existence of a moduli space of G -Higgs principal bundles (for G real algebraic) similar to the one constructed by Simpson for H -Higgs principal bundles (for H complex algebraic). Such a moduli space should exist and enjoy nice properties (this was confirmed to us by Simpson in an e-mail), but we were unable to find such a construction in the literature.

3. THE CASE $G = \text{SU}(p, 2)$

In this section we prove the main theorem for representations into $G = \text{SU}(p, 2)$. However, some arguments and results are valid in the general case $G = \text{SU}(p, q)$, $p \geq q$, and therefore we will specialize to the case $q = 2$ only when necessary.

3.1. The Hermitian symmetric space $\text{SU}(p, q)/\text{S}(\text{U}(p) \times \text{U}(q))$.

3.1.1. General facts.

The reader should consult [He01, Sat80] for details about this section.

Let \mathbb{E} be a complex vector space of dimension $p + q$, with $p \geq q \geq 1$, endowed with a non-degenerate Hermitian form F of signature (p, q) . Let \mathbb{W} be a q -dimensional complex subspace of \mathbb{E} on which F is negative-definite, and let \mathbb{V} be its F -orthogonal complement, so that $\mathbb{E} = \mathbb{V} \oplus \mathbb{W}$. Let $\{e_1, \dots, e_p\}$ and $\{e_{p+1}, \dots, e_{p+q}\}$ be F -orthonormal bases of \mathbb{V} and \mathbb{W} respectively and let $I_{p,q}$ be the matrix of F in the resulting basis of \mathbb{E} .

The symmetric space \mathcal{X} is defined as the space of all q -dimensional complex subspaces of \mathbb{E} on which F is negative-definite. It is an open submanifold of the complex Grassmannian of q -planes of \mathbb{E} . When $q = 1$, \mathcal{X} is complex hyperbolic space of (complex) dimension p which we denote by $\mathbb{H}_{\mathbb{C}}^p$. The group $G = \text{SU}(p, q) = \{g \in \text{SL}(p+q, \mathbb{C}) \mid {}^t \bar{g} I_{p,q} g = I_{p,q}\}$ acts transitively on \mathcal{X} by analytic isomorphisms, while the isotropy subgroup K of G at \mathbb{W} is identified with the maximal compact subgroup $\text{S}(\text{U}(p) \times \text{U}(q))$ of $\text{SU}(p, q)$, so that \mathcal{X} can be identified with

$SU(p, q)/S(U(p) \times U(q))$. The Cartan involution c of G is given by conjugation by the element $\alpha = \xi I_{p, q}$, where $\xi \in \mathbb{C}$ satisfies $\xi^{p+q} = (-1)^q$, so that $\alpha \in SL(p+q, \mathbb{C})$.

Let \mathfrak{g} be the Lie algebra of G , and $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ the Cartan decomposition given by c . $\mathfrak{k} \subset \mathfrak{g}$ is the Lie algebra of K and we have the following matrix expressions:

$$\mathfrak{k} = \left\{ \begin{pmatrix} X_1 & 0 \\ 0 & X_2 \end{pmatrix}, X_1 \in M_p(\mathbb{C}), X_2 \in M_q(\mathbb{C}), {}^t\bar{X}_i = -X_i \ (i = 1, 2), \operatorname{tr} X_1 + \operatorname{tr} X_2 = 0 \right\},$$

$$\mathfrak{p} = \left\{ \begin{pmatrix} 0 & A \\ {}^t\bar{A} & 0 \end{pmatrix}, A \in M_{p, q}(\mathbb{C}) \right\} \simeq_{\mathbb{R}} \operatorname{Hom}_{\mathbb{C}}(\mathbb{W}, \mathbb{V}).$$

The tangent space $T_o\mathcal{X}$ at $o \in \mathcal{X}$ will be identified with \mathfrak{p} . More generally, the tangent bundle $T\mathcal{X}$ of \mathcal{X} is the bundle $G \times_{\operatorname{Ad}K} \mathfrak{p}$ associated to the K -principal bundle $G \rightarrow \mathcal{X} = G/K$ via the adjoint action of K on \mathfrak{p} . The complex structure J on $T_o\mathcal{X}$ is given by

$$J \begin{pmatrix} 0 & A \\ {}^t\bar{A} & 0 \end{pmatrix} = \begin{pmatrix} 0 & \sqrt{-1}A \\ -\sqrt{-1}{}^t\bar{A} & 0 \end{pmatrix}$$

whereas the G -invariant Kähler metric $g_{\mathcal{X}}$ on \mathcal{X} is defined at o by

$$g_{\mathcal{X}}(X, Y) = 2\operatorname{tr}(YX) = 4\operatorname{Re} \operatorname{tr}({}^t\bar{B}A), \text{ if } X = \begin{pmatrix} 0 & A \\ {}^t\bar{A} & 0 \end{pmatrix}, Y = \begin{pmatrix} 0 & B \\ {}^t\bar{B} & 0 \end{pmatrix} \in \mathfrak{p}.$$

The corresponding Kähler form will be denoted by $\omega_{\mathcal{X}} = g_{\mathcal{X}}(J, \cdot)$.

The complexification $G_{\mathbb{C}}$ of G is $SL(p+q, \mathbb{C})$. The complex conjugation of $G_{\mathbb{C}}$ w.r.t. G is given by $\sigma(g) = (I_{p, q} {}^t\bar{g} I_{p, q})^{-1}$, whereas the complex conjugation w.r.t. its compact real form $SU(p+q)$ is given by $\tau(g) = ({}^t\bar{g})^{-1}$. The Lie algebra $\mathfrak{g}_{\mathbb{C}}$ of $G_{\mathbb{C}}$ splits as $\mathfrak{k}_{\mathbb{C}} \oplus \mathfrak{p}_{\mathbb{C}}$ where $\mathfrak{k}_{\mathbb{C}}$ is the Lie algebra of the complexification $K_{\mathbb{C}} = S(\operatorname{GL}(p, \mathbb{C}) \times \operatorname{GL}(q, \mathbb{C}))$ of K and

$$\mathfrak{p}_{\mathbb{C}} = \mathfrak{p} \otimes \mathbb{C} = \left\{ \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}, A \in M_{p, q}(\mathbb{C}), B \in M_{q, p}(\mathbb{C}) \right\} \simeq_{\mathbb{C}} \operatorname{Hom}_{\mathbb{C}}(\mathbb{W}, \mathbb{V}) \oplus \operatorname{Hom}_{\mathbb{C}}(\mathbb{V}, \mathbb{W}).$$

The extended complex structure $J \otimes \operatorname{Id}$ acting on $\mathfrak{p}_{\mathbb{C}}$ has two eigenspaces

$$\mathfrak{p}^{1,0} = \left\{ \begin{pmatrix} 0 & A \\ 0 & 0 \end{pmatrix}, A \in M_{p, q}(\mathbb{C}) \right\} \simeq_{\mathbb{C}} \operatorname{Hom}_{\mathbb{C}}(\mathbb{W}, \mathbb{V})$$

and

$$\mathfrak{p}^{0,1} = \left\{ \begin{pmatrix} 0 & 0 \\ A & 0 \end{pmatrix}, A \in M_{q, p}(\mathbb{C}) \right\} \simeq_{\mathbb{C}} \operatorname{Hom}_{\mathbb{C}}(\mathbb{V}, \mathbb{W}).$$

The complexified tangent bundle $T^{\mathbb{C}}\mathcal{X}$ of \mathcal{X} is isomorphic to $G \times_{\operatorname{Ad}K} \mathfrak{p}_{\mathbb{C}} \simeq (G \times_K K_{\mathbb{C}}) \times_{\operatorname{Ad}K_{\mathbb{C}}} \mathfrak{p}_{\mathbb{C}}$, whereas the holomorphic tangent bundle $T^{1,0}\mathcal{X}$ is isomorphic to $G \times_{\operatorname{Ad}K} \mathfrak{p}^{1,0} \simeq (G \times_K K_{\mathbb{C}}) \times_{\operatorname{Ad}K_{\mathbb{C}}} \mathfrak{p}^{1,0}$. There is a natural Hermitian metric on the holomorphic tangent bundle of \mathcal{X} given on $T_o^{1,0}\mathcal{X} \simeq M_{p, q}(\mathbb{C})$ by $h(A, B) = 4\operatorname{tr}({}^t\bar{B}A)$.

The holomorphic sectional curvature for the complex line $\langle X \rangle$ generated by a nonzero $X = \begin{pmatrix} 0 & A \\ {}^t\bar{A} & 0 \end{pmatrix} \in T_o\mathcal{X}$, or equivalently by a non-zero $\begin{pmatrix} 0 & A \\ 0 & 0 \end{pmatrix} \in T_o^{1,0}\mathcal{X}$, is given by

$$K(\langle X \rangle) = -\frac{\operatorname{tr}(({}^t\bar{A}A)^2)}{(\operatorname{tr}({}^t\bar{A}A))^2}.$$

This formula shows that $K(\langle X \rangle)$ is pinched between -1 and $-1/q$ and that $K(\langle X \rangle) = -1/q$ if and only if the column vectors of A are pairwise orthogonal and have the same norm (for the standard Hermitian scalar product in \mathbb{C}^p).

The metric $g_{\mathcal{X}}$ is Einstein and with our normalization, its Ricci curvature tensor is $-\frac{p+q}{2} g_{\mathcal{X}}$.

3.1.2. Maximal embeddings.

There is a natural identification of $\mathcal{X} = \mathrm{SU}(p, q)/\mathrm{S}(\mathrm{U}(p) \times \mathrm{U}(q))$ with the space $\{Z \in M_{p, q}(\mathbb{C}), I_q - {}^t Z Z > 0\}$ [Sat80]. Therefore if $m \leq p/q$, we have a holomorphic totally geodesic embedding of complex hyperbolic space $\mathbb{H}_{\mathbb{C}}^m = \mathrm{SU}(m, 1)/\mathrm{S}(\mathrm{U}(m) \times \mathrm{U}(1))$ into \mathcal{X} given by

$$(\star) \quad \mathbb{H}_{\mathbb{C}}^m \ni z = \begin{pmatrix} z_1 \\ z_2 \\ \vdots \\ z_m \end{pmatrix} \mapsto Z = \begin{pmatrix} z_1 I_q \\ z_2 I_q \\ \vdots \\ z_m I_q \\ 0_{p-qm, q} \end{pmatrix} \in \mathcal{X}.$$

This shows that \mathcal{X} contains totally geodesic copies of complex hyperbolic m -space of holomorphic sectional curvature $-1/q$, for all $m \leq p/q$. The next lemma implies that $[p/q]$ is the maximal possible dimension of such submanifolds:

Lemma 3.1. *Let S be a complex linear subspace of $T_o^{1,0} \mathcal{X}$. If, for every nonzero $A \in S$, $K(\langle A \rangle) = -1/q$, then $\dim_{\mathbb{C}} S \leq p/q$.*

Proof. The metric and the holomorphic sectional curvature are obviously invariant under the left action of $\mathrm{U}(p)$ on $T_o^{1,0} \mathcal{X} \simeq M_{p, q}(\mathbb{C})$.

Let $\{A_1, \dots, A_d\}$ be an orthonormal basis of S . We are going to show that there exists $U \in \mathrm{U}(p)$ such that

$$U A_k = \frac{1}{2\sqrt{q}} \begin{pmatrix} 0_{(k-1)q, q} \\ I_q \\ 0_{p-kq, q} \end{pmatrix}, \quad k = 1, \dots, d.$$

Since the column vectors of each A_i are pairwise orthogonal and have the same norm, there exists $U_1 \in \mathrm{U}(p)$ such that

$$U_1 A_1 = \frac{1}{2\sqrt{q}} \begin{pmatrix} I_q \\ 0_{p-q, q} \end{pmatrix}.$$

Now, for any $(\lambda, \mu) \in \mathbb{C}^2 \setminus \{(0, 0)\}$, ${}^t(\lambda \bar{A}_1 + \mu \bar{A}_2)(\lambda A_1 + \mu A_2)$ must be a (nonzero) multiple of I_q . This implies that for any (λ, μ) , $\lambda \bar{\mu} {}^t \bar{A}_1 A_2 + \lambda \bar{\lambda} A_1 {}^t \bar{A}_2$ is a multiple of I_q (because ${}^t \bar{A}_1 A_1$ and ${}^t \bar{A}_2 A_2$ are). Moreover, it is trace free because A_1 and A_2 are orthogonal. Thus, ${}^t \bar{A}_1 A_2 = 0$ (that is each column vector of A_1 is orthogonal to every column vector of A_2) and there exists $U_2 \in \mathrm{U}(p)$ such that $U_2 U_1 A_1 = U_1 A_1$ and

$$U_2 U_1 A_2 = \frac{1}{2\sqrt{q}} \begin{pmatrix} 0_{q, q} \\ I_q \\ 0_{p-2q, q} \end{pmatrix}.$$

One might continue this process and after d steps, one obtains $U = U_d \dots U_1$. It is then clear that d must be less than or equal to p/q . \square

The embedding (\star) will be denoted by f_{max} and called the maximal embedding of $\mathbb{H}_{\mathbb{C}}^m$ into \mathcal{X} . This is because $f_{max}^* g_{\mathcal{X}} = qg$, that is, for any $x \in \mathbb{H}_{\mathbb{C}}^m$ and any $X \in T_x \mathbb{H}_{\mathbb{C}}^m$, $g_{\mathcal{X}}(\mathrm{d}f_{max}(X), \mathrm{d}f_{max}(X)) = qg(X, X)$, while for a general holomorphic map $f : \mathbb{H}_{\mathbb{C}}^m \rightarrow \mathcal{X}$ we only know that $f^* g_{\mathcal{X}} \leq qg$ from the Ahlfors-Schwarz-Pick lemma (see for example [Ro80, Theorem 2]). Moreover,

Proposition 3.2. *Let $f : \mathbb{H}_{\mathbb{C}}^m \longrightarrow \mathcal{X} = \mathrm{SU}(p, q)/\mathrm{S}(\mathrm{U}(p) \times \mathrm{U}(q))$ be a holomorphic map such that $f^*g_{\mathcal{X}} = qg$ holds everywhere. Then $p/q \geq m$ and f is totally geodesic. In fact, up to composition with an isometry of \mathcal{X} , f is the maximal embedding f_{max} .*

Proof. If X is a nonzero tangent vector at $x \in \mathbb{H}_{\mathbb{C}}^m$, we denote by $C_{x, X}$ the complex geodesic through x that is tangent to X . Let z be a (global) complex coordinate on $C_{x, X}$ and let $\Sigma^2 |dz|^2$ (resp. $\sigma^2 |dz|^2$) be the Hermitian metric induced by g (resp. $f^*g_{\mathcal{X}} = qg$) on $C_{x, X}$. The Gaussian curvature of Σ (resp. σ) is given by $K = -\frac{1}{\Sigma^2} \Delta \log \Sigma$ (resp. $k = -\frac{1}{\sigma^2} \Delta \log \sigma$). Since $C_{x, X} \subset \mathbb{H}_{\mathbb{C}}^m$ is totally geodesic, we have $K \equiv -1$. Moreover, because of the holomorphicity of f , k is bounded from above by $-1/q$ which is the maximum of the holomorphic sectional curvature on \mathcal{X} and $k \equiv -1/q$ iff the restriction of f to $C_{x, X}$ is totally geodesic. But $\sigma = \sqrt{q} \Sigma$ and so

$$k = -\frac{1}{\sigma^2} \Delta \log \sigma = -\frac{1}{q\Sigma^2} \Delta \log \Sigma = -\frac{1}{q}.$$

Thus, $k \equiv -1/q$ and, since this is true for any (x, X) , f must be totally geodesic.

Let $o = 0_{m,1}$ (resp. $o' = 0_{p,q}$) be fixed origins in $\mathbb{H}_{\mathbb{C}}^m$ (resp. in \mathcal{X}). One may suppose (after composition with an isometry of \mathcal{X}) that $f(o) = o'$. A consequence of the preceding discussion is that $df(T_o\mathbb{H}_{\mathbb{C}}^m)$ is an m -dimensional complex subspace of $T_{o'}\mathcal{X}$ on which the restriction of the holomorphic sectional curvature is constant, equal to $-1/q$. By the proof of Lemma 3.1, we know that $m \geq p/q$ and that, after composition of f with a suitable isometry, $df|_{T_o\mathbb{H}_{\mathbb{C}}^m} = df_{max}|_{T_o\mathbb{H}_{\mathbb{C}}^m}$. By uniqueness of the totally geodesic map satisfying this condition, one has $f = f_{max}$. \square

Another maximal embedding of $\mathbb{H}_{\mathbb{C}}^m$ into $\mathcal{X} = \mathrm{SU}(p, q)/\mathrm{S}(\mathrm{U}(p) \times \mathrm{U}(q))$ ($p/q \geq m$) is given by

$$f'_{max} : z = \begin{pmatrix} z_1 \\ z_2 \\ \vdots \\ z_m \end{pmatrix} \longmapsto Z = \begin{pmatrix} z & 0 & \cdots & 0 \\ 0 & z & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & z \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 \end{pmatrix}.$$

From the previous proposition (it can also be easily verified by hand), this embedding is equal to f_{max} composed with an isometry of \mathcal{X} . The geometric picture is maybe clearer here: f'_{max} is a diagonal embedding of $\mathbb{H}_{\mathbb{C}}^m$ into $(\mathbb{H}_{\mathbb{C}}^m)^q \subset \mathcal{X}$ corresponding to a (diagonal) embedding of $\mathrm{SU}(m, 1)$ into $\mathrm{SU}(m, 1)^q \subset \mathrm{SU}(p, q)$. Moreover the stabilizer in $\mathrm{SU}(p, q)$ of the image of f'_{max} can be computed quite easily. First, let us consider the subgroup of $\mathrm{U}(p, q)$ consisting of elements of the form:

$$\begin{pmatrix} A_{11} & \cdots & A_{1q} & 0 & B_{11} & \cdots & B_{1q} \\ \vdots & \cdots & \vdots & 0 & \vdots & \cdots & \vdots \\ A_{q1} & \cdots & A_{qq} & 0 & B_{q1} & \cdots & B_{qq} \\ 0 & \cdots & 0 & U & 0 & \cdots & 0 \\ C_{11} & \cdots & C_{1q} & 0 & d_{11} & \cdots & d_{1q} \\ \vdots & \cdots & \vdots & 0 & \vdots & \cdots & \vdots \\ C_{q1} & \cdots & C_{qq} & 0 & d_{q1} & \cdots & d_{qq} \end{pmatrix}$$

where $A_{ij} \in M_m(\mathbb{C})$, $B_{ij} \in M_{m,1}(\mathbb{C})$, $C_{ij} \in M_{1,m}(\mathbb{C})$, $d_{ij} \in \mathbb{C}$ and $U \in \mathrm{U}(r)$ ($r = p - qm$).

Let us denote by S_q the symmetric group on q letters, and by $\mathrm{U}(1)^q \rtimes S_q$ the semi-direct product of $\mathrm{U}(1)^q$ by S_q given by the group operation $(\alpha, \sigma) \cdot (\beta, \tau) = (\alpha \cdot \sigma(\beta), \tau \circ \sigma)$.

Define a group homomorphism φ of $(U(1)^q \times S_q) \times SU(m, 1) \times U(r)$ in the above subgroup of $U(p, q)$ in the following way: if $\alpha = (\alpha_1, \dots, \alpha_q) \in U(1)^q$, $\sigma \in S_q$, $u \in U(r)$ and

$$g = \begin{pmatrix} A & B \\ C & d \end{pmatrix} \in SU(m, 1) \quad (\text{where } A \in M_m(\mathbb{C}), B \in M_{m,1}(\mathbb{C}), C \in M_{1,m}(\mathbb{C}), d \in \mathbb{C}),$$

the image of (α, σ, g, u) in $U(p, q)$ is the matrix defined by $A_{i\sigma(i)} = \alpha_i A$, $B_{i\sigma(i)} = \alpha_i B$, $C_{i\sigma(i)} = \alpha_i C$, $d_{i\sigma(i)} = \alpha_i d$, $U = u$, and the other coefficients are zero.

Then $\text{Ker}\varphi$ is isomorphic to $\mathbb{Z}/(m+1)\mathbb{Z}$ and the stabilizer in $SU(p, q)$ of the image of f'_{max} is $\text{Im}\varphi \cap SU(p, q)$.

3.2. Toledo invariant and $SU(p, 2)$ -Higgs bundles.

Now we consider a reductive representation ρ of a torsion-free uniform lattice of $SU(m, 1)$, $m > 1$, into the Lie group of Hermitian type $G = SU(p, q)$, $p \geq q \geq 1$. Let M be the closed complex hyperbolic manifold $\Gamma \backslash \mathbb{H}_{\mathbb{C}}^m$. As explained in the introduction, the Toledo invariant can be expressed using the degree of the pull-back of the holomorphic tangent bundle $T^{1,0}\mathcal{X}$ of $\mathcal{X} = G/K$ by any ρ -equivariant map $f : \mathbb{H}_{\mathbb{C}}^m \rightarrow \mathcal{X}$, which we can choose to be harmonic. Let then $(P_{K_{\mathbb{C}}}, \theta)$ be the G -Higgs principal bundle on M associated to ρ and f as in section 2 and let E be the holomorphic vector bundle on M associated to $P_{K_{\mathbb{C}}}$ via the action of $K_{\mathbb{C}}$ on $\mathbb{E} = \mathbb{V} \oplus \mathbb{W}$. Since $K_{\mathbb{C}}$ respects the decomposition $\mathbb{E} = \mathbb{V} \oplus \mathbb{W}$, the bundle E splits holomorphically as the sum of the rank p subbundle $V = P_{K_{\mathbb{C}}} \times_{K_{\mathbb{C}}} \mathbb{V}$ with the rank q subbundle $W = P_{K_{\mathbb{C}}} \times_{K_{\mathbb{C}}} \mathbb{W}$. As a differentiable bundle E is the bundle associated to the flat principal bundle P_G via the action of G on \mathbb{E} : it is flat, hence of degree 0. In particular, $\deg V = -\deg W$.

The Higgs field θ is a holomorphic $(1,0)$ -form taking values in the bundle $P_{K_{\mathbb{C}}} \times_{\text{Ad}K_{\mathbb{C}}} \mathfrak{p}_{\mathbb{C}} = \text{Hom}(W, V) \oplus \text{Hom}(V, W)$ so that we can write (see also [Xi00, BGG03])

$$\theta = \begin{pmatrix} 0 & \beta \\ \gamma & 0 \end{pmatrix}, \quad \text{with } \begin{cases} \beta : W \rightarrow V \otimes \Omega^1 \\ \gamma : V \rightarrow W \otimes \Omega^1 \end{cases}$$

where Ω^1 is the sheaf (of germs) of holomorphic 1-forms on M . An important fact that we shall use is that Ω^1 and its dual T^1 are semistable sheaves in the usual sense (in fact they are stable), because M is complex hyperbolic, hence Kähler-Einstein. Recall that the Higgs vector bundle (E, θ) is polystable (as a Higgs bundle).

It is clear that the bundle $f^*T^{1,0}\mathcal{X}$ is nothing but the bundle $\text{Hom}(W, V)$ and therefore its degree is simply given by $p \deg W^* + q \deg V = -(p+q)\deg W$. We obtain that $\tau(\rho) = \frac{4\pi}{m!} \deg W$, so that the Milnor-Wood type inequality reads

$$|\deg W| \leq \frac{q}{m+1} \deg \Omega^1.$$

Therefore, for $q = 2$ and ρ reductive, our main theorem can be reformulated:

Theorem 3.3. *Let Γ be a torsion free uniform lattice in $SU(m, 1)$, $m > 1$, and let $\rho : \Gamma \rightarrow SU(p, 2)$ be a reductive representation. Let $E = V \oplus W$ be the $SU(p, 2)$ -Higgs vector bundle on $M = \Gamma \backslash \mathbb{H}_{\mathbb{C}}^m$ associated to ρ . Then $|\deg W| \leq \frac{2}{m+1} \deg \Omega^1$ with equality if and only if $m \leq p/2$ and, up to conjugacy, ρ is induced by the maximal embedding $f_{max} : \mathbb{H}_{\mathbb{C}}^m \rightarrow \mathcal{X}$ or by its conjugate.*

If we deform the Higgs bundle (E, θ) via the \mathbb{C}^* -action on the moduli space as in section 2, we obtain a system of Hodge bundles:

$$E_1 \xrightarrow{\theta_1} E_2 \xrightarrow{\theta_2} \dots \xrightarrow{\theta_{k-1}} E_k \xrightarrow{\theta_k} 0.$$

Moreover, each subbundle E_i splits as $E_i = V_i \oplus W_i$ with $V_i \subset V$ and $W_i \subset W$, and θ_i decomposes as $\gamma_i \oplus \beta_i$, where $\gamma_i : V_i \rightarrow W_{i+1} \otimes \Omega^1$ and $\beta_i : W_i \rightarrow V_{i+1} \otimes \Omega^1$. We obtain two Higgs subbundles

$$V_1 \rightarrow W_2 \rightarrow V_3 \rightarrow W_4 \rightarrow \dots \rightarrow 0$$

and

$$W_1 \rightarrow V_2 \rightarrow W_3 \rightarrow V_4 \rightarrow \dots \rightarrow 0$$

which are again polystable of degree 0.

So far everything we said was valid in the general rank q case. Now we will need the assumption that $q = 2$ to ensure that the systems of Hodge bundles we obtain are simple ones and/or that the decomposition of W is maximal in the sense that W splits into a sum of line bundles. Indeed, if W has rank 2, we see that by regrouping and renaming the subbundles if necessary, we can write our system of Hodge bundles either as a polystable Higgs bundle of the form

$$V_1 \xrightarrow{\gamma_1} W \xrightarrow{\beta} V_2 \xrightarrow{\gamma_2} 0$$

with $V_1 \oplus V_2 = V$, or as a polystable Higgs bundle of the form

$$V_1 \xrightarrow{\gamma_1} W_1 \xrightarrow{\beta_1} V_2 \xrightarrow{\gamma_2} W_2 \xrightarrow{\beta_2} V_3 \xrightarrow{\gamma_3} 0$$

where W_1 and W_2 are line bundles, $W_1 \oplus W_2 = W$ and $V_1 \oplus V_2 \oplus V_3 = V$.

3.3. Proof of the Milnor-Wood type inequality.

The case of non reductive representations will be postponed to Paragraph 3.3.3. Until then, the representation $\rho : \Gamma \rightarrow \mathrm{SU}(p, 2)$ is assumed to be reductive so that we can apply the results of Section 3.2. Our proof of the inequality $|\mathrm{deg} W| \leq \frac{2}{m+1} \mathrm{deg} \Omega^1$ will be different according to the form of the system of Hodge bundles we obtain by deforming the polystable Higgs bundles $E = V \oplus W$ via the \mathbb{C}^* -action. Note that the deformation changes the holomorphic structures of E , V and W , but not their isomorphism classes as differentiable complex vector bundles, hence their degrees remain unchanged. During the proof, we will see that if equality holds, some bundles have stability properties (in the usual sense) that will be useful for the study of maximal representations in Section 3.4.

We refer to [VZ07, HL97] for general facts about sheaves and stability.

3.3.1. System of Hodge bundles of the type $V_1 \rightarrow W \rightarrow V_2 \rightarrow 0$.

Here the important point is that the system of Hodge bundles we are dealing with is a ternary bundle, and no limitation on the rank of W is needed. Hence the results of this paragraph are valid in the general case $\mathrm{rk} W = q \geq 1$.

Let \mathcal{F} be the maximal destabilizing subsheaf of W , that is, the first term in the Harder-Narasimhan filtration of W [VZ07, HL97]. By definition, \mathcal{F} has maximal slope among the subsheaves of W , hence is semistable. Consider the restriction $\beta_{\mathcal{F}} : \mathcal{F} \otimes T^1 \rightarrow V_2$. Since θ vanishes on V_2 , $\mathcal{F} \oplus \mathrm{Im} \beta_{\mathcal{F}}$ is a Higgs subsheaf and hence by stability, $\mathrm{deg} \mathrm{Im} \beta_{\mathcal{F}} \leq -\mathrm{deg} \mathcal{F}$. Now, the tensor product of two semistable sheaves is again semistable and hence $\mathcal{F} \otimes T^1$ is semistable. Therefore we have $\mu(\mathcal{F}) + \mu(T^1) = \mu(\mathcal{F} \otimes T^1) \leq \mu(\mathrm{Im} \beta_{\mathcal{F}})$ which implies $(\mathrm{rk} \beta_{\mathcal{F}} + \mathrm{rk} \mathcal{F})\mu(\mathcal{F}) \leq \mathrm{rk} \beta_{\mathcal{F}} \mu(\Omega^1)$. Thus,

$$\mathrm{deg} W \leq q\mu(\mathcal{F}) \leq q \frac{\mathrm{rk} \beta_{\mathcal{F}}}{\mathrm{rk} \beta_{\mathcal{F}} + \mathrm{rk} \mathcal{F}} \frac{\mathrm{deg} \Omega^1}{m} \leq \frac{q}{m+1} \mathrm{deg} \Omega^1$$

where the first inequality follows from the fact that \mathcal{F} is of maximal slope among the subsheaves in W , and the last from $\mathrm{rk} \beta_{\mathcal{F}} \leq m \mathrm{rk} \mathcal{F}$.

The inequality $\deg W \geq -\frac{q}{m+1}\deg \Omega^1$ is obtained exactly in the same way by considering the dual Higgs bundle

$$V_2^* \xrightarrow{t\beta} W^* \xrightarrow{t\gamma_1} V_1^* \longrightarrow 0.$$

Assume that equality holds, for example that $\deg W = \frac{q}{m+1}\deg \Omega^1$. Then, retracing our steps, we see that W must be a semistable bundle (in the usual sense), that we must have $\text{rk } \beta = mq$, i.e. $\beta : W \otimes T^1 \rightarrow V$ generically injective, and moreover that $\deg W \oplus \text{Im } \beta = 0$. This last fact implies by polystability that E splits as the sum of $(W \oplus \text{Im } \beta, \beta)$ with another polystable Higgs bundle E' of degree 0. In our case, this means that V_2 splits holomorphically as $\text{Im } \beta \oplus V_2'$ and that γ_1 vanishes. E' is then the polystable (in the usual sense) subbundle $V_1 \oplus V_2'$ of V .

In the same manner, we find that $\deg V = \frac{q}{m+1}\deg \Omega^1$ implies that W is a semistable bundle, $\beta = 0$, V_1 splits holomorphically as $\text{Ker } \gamma_1 \oplus V_1'$, and $\gamma_1 : V_1' \rightarrow W \otimes \Omega^1$ is an isomorphism. Our system of Hodge bundles is the sum of the Higgs bundle $(V_1' \oplus W, \gamma_1)$ with the polystable subbundle $\text{Ker } \gamma_1 \oplus V_2$ of V .

3.3.2. System of Hodge bundles of the type $V_1 \rightarrow W_1 \rightarrow V_2 \rightarrow W_2 \rightarrow V_3 \rightarrow 0$.

Here, we need to assume that $\text{rk } W = q = 2$, namely that W_1 and W_2 are line bundles.

Assume first that $\beta_1 : W_1 \otimes T^1 \rightarrow V_2$ vanishes. We then have to deal with the sum of two polystable Higgs bundles of degree 0: $V_1 \rightarrow W_1 \rightarrow 0$ and $V_2 \rightarrow W_2 \rightarrow V_3 \rightarrow 0$. In this situation, we already know from 3.3.1 that $0 \leq \deg V_1 \leq \frac{1}{m+1}\deg \Omega^1$ and $|\deg(V_2 \oplus V_3)| \leq \frac{1}{m+1}\deg \Omega^1$ which gives the result.

We are left with the case where $\beta_1 : W_1 \otimes T^1 \rightarrow V_2$ is non zero.

Consider $\gamma_1 : V_1 \rightarrow W_1 \otimes \Omega^1$. We have $\deg V_1 = \deg \text{Ker } \gamma_1 + \deg \text{Im } \gamma_1 \leq \text{rk } \gamma_1 \mu(W_1 \otimes \Omega^1)$ since $\text{Ker } \gamma_1$ is θ -invariant and $W_1 \otimes \Omega^1$ is a semistable bundle, being the product of a stable bundle by a line bundle. Since $\deg W_1 = -(\deg V_1 + \deg V_2 + \deg W_2 + \deg V_3)$ we obtain

$$\deg V_1 \leq \frac{\text{rk } \gamma_1}{1 + \text{rk } \gamma_1} \frac{\deg \Omega^1}{m} - \frac{\text{rk } \gamma_1}{1 + \text{rk } \gamma_1} (\deg V_2 + \deg W_2 + \deg V_3).$$

In the same way,

$$\deg V_2 \leq \frac{\text{rk } \gamma_2}{1 + \text{rk } \gamma_2} \frac{\deg \Omega^1}{m} - \frac{\text{rk } \gamma_2}{1 + \text{rk } \gamma_2} (\deg V_1 + \deg W_1 + \deg V_3).$$

Hence,

$$\begin{aligned} \deg V \leq & \left(\frac{\text{rk } \gamma_1}{1 + \text{rk } \gamma_1} + \frac{\text{rk } \gamma_2}{1 + \text{rk } \gamma_2} \right) \frac{\deg \Omega^1}{m} + \\ & \left(\frac{\text{rk } \gamma_2}{1 + \text{rk } \gamma_2} - \frac{\text{rk } \gamma_1}{1 + \text{rk } \gamma_1} \right) (\deg V_2 + \deg W_2 + \deg V_3) + \left(1 - \frac{\text{rk } \gamma_2}{1 + \text{rk } \gamma_2} \right) \deg V_3. \end{aligned}$$

Now the commutation relation $[\theta, \theta] = 0$ gives us control on the rank of the γ_i 's:

Lemma 3.4. *Assume that $\beta_i : W_i \otimes T^1 \rightarrow V_{i+1}$ is non zero. Then the rank of $\gamma_i : V_i \rightarrow W_i \otimes \Omega^1$ is at most 1.*

Proof. This is linear algebra. We work in a single fiber above some point in M . We write $W_i = \mathbb{C}w_i$. Since β_i is non zero, there exists $Z \in T^1$ such that $\beta_i(Z)w_i \neq 0$. Assume that the rank of γ_i is at least 2 at some point. Then we can find two linearly independent forms α, α' and two vectors v, v' in V_i such that $\gamma_i(v) = w_i \otimes \alpha$ and $\gamma_i(v') = w_i \otimes \alpha'$. Now, the commutation relation $[\theta, \theta] = 0$ means in particular that for all $X, Y \in T^1$ and all $u \in V$, $\beta_i(X)\gamma_i(Y)u = \beta_i(Y)\gamma_i(X)u$.

If we take $u = v$, $Y = Z$ and X such that $\alpha(X) \neq 0$, we get $\alpha(Z)\beta_i(X)w_i = \alpha(X)\beta_i(Z)w_i$, which implies that $\alpha(Z)$ and $\beta_i(X)w_i$ are different from zero. So if now X' is such that $\alpha(X') = 0$, $\beta_i(X')w_i = 0$. We can choose such an X' with the additional property that $\alpha'(X') \neq 0$, since α and α' are independent. This is a contradiction since we could have taken $u = v'$ to prove that $\beta_i(X')w_i \neq 0$ if $\alpha'(X') \neq 0$. \square

We assumed that $\beta_1 : W_1 \otimes T^1 \rightarrow V_2$ is non zero, hence $\text{rk } \gamma_1 \leq 1$. If $\text{rk } \gamma_2 \geq \text{rk } \gamma_1$, using the fact that $\deg V_2 + \deg W_2 + \deg V_3$ and $\deg V_3$ are both non positive, and since $\text{rk } \gamma_2 \leq m$, we get

$$\deg V \leq \frac{3m+1}{2m(m+1)} \deg \Omega^1 \leq \frac{2}{m+1} \deg \Omega^1,$$

and the last inequality is strict as soon as $m > 1$. If $\text{rk } \gamma_2 < \text{rk } \gamma_1$, that is if $\text{rk } \gamma_1 = 1$ and $\text{rk } \gamma_2 = 0$, we are in the case $V_1 \rightarrow W_1 \rightarrow V_2 \rightarrow 0$ and hence

$$\deg V \leq \deg(V_1 \oplus V_2) = -\deg W_1 \leq \frac{1}{m+1} \deg \Omega^1.$$

Again, the lower bound on $\deg V$ is obtained by considering the dual Higgs bundle

$$V_3^* \xrightarrow{t\beta_2} W_2^* \xrightarrow{t\gamma_2} V_2^* \xrightarrow{t\beta_1} W_1^* \xrightarrow{t\gamma_1} V_1^* \rightarrow 0.$$

Assume we are in the equality case and $m > 1$. If $\deg V = \frac{2}{m+1} \deg \Omega^1$, $\beta_1 : W_1 \otimes T^1 \rightarrow V_2$ vanishes, and the equalities $\deg V_1 = \frac{1}{m+1} \deg \Omega^1 = \deg(V_2 \oplus V_3)$ hold. We saw in 3.3.1 that in this situation, $\beta_2 = 0$ and there exists holomorphic subbundles $V'_i \subset V_i$ such that $\gamma_i : V'_i \rightarrow W_i \otimes \Omega^1$ are isomorphisms for $i = 1, 2$.

If $\deg W = \frac{2}{m+1} \deg \Omega^1$, we find that for $i = 1, 2$, $\gamma_i = 0$ and there exists a holomorphic subbundle $V'_i \subset V_i$ such that $\beta_i : W_i \otimes T^1 \rightarrow V'_i$ is an isomorphism.

In either cases, $\deg W_1 = \deg W_2$ and W with the deformed complex structure is polystable hence semistable.

3.3.3. Non reductive representations.

Assume now that the representation $\rho : \Gamma \rightarrow G = \text{SU}(p, 2)$ is not reductive. This implies that $\rho(\Gamma)$ fixes a point ξ on the boundary at infinity $\mathcal{X}(\infty)$ of \mathcal{X} ([La91]). Let us fix an origin $o \in \mathcal{X}$ and let c be the unit speed geodesic ray starting from o representing ξ . Let $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ be the Cartan decomposition of \mathfrak{g} associated to o and let $X \in \mathfrak{p}$ be such that $X = \dot{c}(0)$ in the usual identification of \mathfrak{p} with $T_o\mathcal{X}$. We have the following description of the stabilizer G_ξ of ξ in G (see for example [Eb96]):

$$G_\xi = \{g \in G \mid \lim_{t \rightarrow +\infty} \exp(-tX) g \exp(tX) \text{ exists}\} = K_\xi \cdot A_\xi \cdot N_\xi$$

where $A_\xi = \exp(\{Y \in \mathfrak{p} \mid [X, Y] = 0\})$, $N_\xi = \{g \in G_\xi \mid \lim_{t \rightarrow +\infty} \exp(-tX) g \exp(tX) = 1\}$, and $K_\xi = G_\xi \cap K$.

By assumption $\rho(\Gamma) \subset G_\xi$ and we can consider the so-called semi-simplification ρ_{ss} of ρ which is defined by $\rho_{ss}(\gamma) = \lim_{t \rightarrow +\infty} \exp(-tX) \rho(\gamma) \exp(tX) \in K_\xi \cdot A_\xi$ for all $\gamma \in \Gamma$. The representation ρ_{ss} belongs to the connected component of ρ in the space $\text{Hom}(\Gamma, G)$ and is reductive: we can apply the results of the last paragraphs to get the Milnor-Wood bound on $\tau(\rho) = \tau(\rho_{ss})$.

In fact we can do better. The representation ρ_{ss} stabilizes the orbit $K_\xi \cdot A_\xi \cdot o = A_\xi \cdot o$, which is a totally geodesic submanifold of \mathcal{X} . It is not difficult to see (and probably well known) that this orbit is either a totally real totally geodesic submanifold of \mathcal{X} (for example, if c is a regular geodesic, it is the unique maximal flat, isometric to \mathbb{R}^2 in our case, containing c), or the Riemannian product of \mathbb{R} with a totally geodesic copy of complex hyperbolic space

$\mathbb{H}_{\mathbb{C}}^{p-1}$ (of induced holomorphic sectional curvature -1). In the first case the Toledo invariant is zero since the restriction of the Kähler form $\omega_{\mathcal{X}}$ to a totally real submanifold vanishes. In the second one it is bounded (in absolute value) by $\text{Vol}(M)$.

Therefore non reductive representations can not be maximal.

3.4. Maximal representations.

Thanks to the previous paragraph, we know that if the representation ρ is maximal, it is reductive. Therefore we may consider the polystable Higgs bundles $(E = V \oplus W, \theta)$ associated to ρ .

In order to prove the Milnor-Wood type inequality $|\text{deg } W| \leq \frac{2}{m+1} \text{deg } \Omega^1$, we deformed the Higgs bundle E to a system of Hodge bundles. Here, we need to distinguish between these two Higgs bundles, and we will call the latter $(E_0 = V_0 \oplus W_0, \theta_0)$. Let $\bar{\partial}_W$ and $\bar{\partial}_{W_0}$ be the complex structure of W and W_0 . Again, although the complex structure is (by definition) not modified by the \mathbb{C}^* -action, in the limit $\bar{\partial}_{W_0}$ is a priori different from $\bar{\partial}_W$. In fact, all we know is that there exist C^∞ -automorphisms g_{t_i} of E such that $g_{t_i}^* \bar{\partial}_W$ goes to $\bar{\partial}_{W_0}$ when t_i goes to 0. Let us call W_{t_i} the bundle W with the complex structure $g_{t_i}^* \bar{\partial}_W$.

The main point of Section 3.3, apart from the proof of the inequality itself, was that $|\text{deg } W| = \frac{2}{m+1} \text{deg } \Omega^1$ implies that the bundle W_0 is semistable (in the usual sense). This implies that W itself is semistable (regardless of the rank of W):

Lemma 3.5. *Assume that W with its initial complex structure $\bar{\partial}_W$ is not a semistable bundle. Then W_0 , that is W with the complex structure $\bar{\partial}_{W_0}$, is not semistable either.*

Proof. Let \mathcal{F} be a subsheaf of $(W, \bar{\partial}_W)$ such that $\mu(\mathcal{F}) > \mu(W)$. Let r be the rank of \mathcal{F} . We have a monomorphism of sheaves $\mathcal{F} \rightarrow W$ and therefore, for all $i \in \mathbb{N}$, we obtain a monomorphism of sheaves $\mathcal{F} \rightarrow W_{t_i}$. This gives non trivial holomorphic maps between the determinant bundle $\det \mathcal{F} = (\bigwedge^r \mathcal{F})^{**}$ of \mathcal{F} and $\bigwedge^r W_{t_i}$. This means that for all $i \in \mathbb{N}$, the cohomology group $H^0(M, \text{Hom}(\det \mathcal{F}, \bigwedge^r W_{t_i}))$ is at least one dimensional. By the upper semicontinuity of cohomology (see Kobayashi [Ko87]), there exists a non-trivial holomorphic map $\det \mathcal{F} \rightarrow \bigwedge^r W_0$. Let I be its image. Since $\det \mathcal{F}$ is stable (it is a line bundle), we have

$$\mu(I) \geq \mu(\det \mathcal{F}) = r\mu(\mathcal{F}) > r\mu(W) = r\mu(W_0) = \mu\left(\bigwedge^r W_0\right).$$

Hence $\bigwedge^r W_0$ is not semistable, and neither is W_0 . \square

Summing up, we proved

Proposition 3.6. *Let $\rho : \Gamma \rightarrow \text{SU}(p, 2)$ be a maximal representation and let $E = V \oplus W$ be the associated $\text{SU}(p, 2)$ -Higgs bundle on $M = \Gamma \backslash \mathbb{H}_{\mathbb{C}}^m$. Then W is a semistable holomorphic bundle.*

The semistability of W is a very strong property and Theorem 3.3 follows from

Theorem 3.7. *Let Γ be a torsion-free uniform lattice in $\text{SU}(m, 1)$, $m > 1$, and $\rho : \Gamma \rightarrow \text{SU}(p, q)$, $p \geq q \geq 1$, be a reductive representation. Let $E = V \oplus W$ be the associated $\text{SU}(p, q)$ -Higgs bundle on $M = \Gamma \backslash \mathbb{H}_{\mathbb{C}}^m$. Assume moreover that W is semistable. Then $|\text{deg } W| \leq \frac{q}{m+1} \text{deg } \Omega^1$, with equality if and only if $m \leq p/q$ and, up to conjugacy, ρ is induced by the maximal embedding $f_{\max} : \mathbb{H}_{\mathbb{C}}^m \rightarrow \mathcal{X}$ or by its conjugate.*

Proof. Consider $\beta : W \otimes T^1 \rightarrow V$ and argue as in 3.3.1 with $\mathcal{F} = W$ and $V_2 = V$ to get the bound $\text{deg } W \leq \frac{q}{m+1} \text{deg } \Omega^1$, with equality if β is injective and $W \oplus \text{Im } \beta$ has degree zero and hence is a polystable Higgs subbundle of E . Now, we have the

Lemma 3.8. *If $m > 1$ and $\beta : W \otimes T^1 \rightarrow V$ is injective, γ vanishes identically.*

Proof of the lemma. This is again a consequence of the relation $[\theta, \theta] = 0$, which in our case reads $\beta(X)\gamma(Y)v = \beta(Y)\gamma(X)v$ for all $X, Y \in T^1$ and all $v \in V$. Let $\{w_1, \dots, w_q\}$ be a basis of W above some point $x \in M$. We can write $\gamma(X)v = \sum_{i=1}^q \lambda_i(X, v)w_i$ and $\gamma(Y)v = \sum_{i=1}^q \lambda_i(Y, v)w_i$. But this implies that

$$\sum_{i=1}^q \lambda_i(Y, v)\beta(X)w_i = \sum_{i=1}^q \lambda_i(X, v)\beta(Y)w_i.$$

Since $m > 1$, we can take X and Y to be linearly independent and the injectivity of β implies that $\beta(X)w_1, \dots, \beta(X)w_q, \beta(Y)w_1, \dots, \beta(Y)w_q$ are linearly independent vectors in V , hence that $\lambda_i(X, v) = \lambda_i(Y, v) = 0$ for all i , namely that $\gamma = 0$. \square

Therefore $\deg W = \frac{q}{m+1} \deg \Omega^1$ implies that $\gamma = 0$, hence that $\partial^{0,1}f = 0$, i.e. the harmonic map f is holomorphic. The theorem easily follows. We know from the Ahlfors-Schwarz-Pick lemma that $f^*g_{\mathcal{X}} \leq qg$. But this implies that the inequality $\langle f^*\omega_{\mathcal{X}}, \omega \rangle \leq 2mq$ is pointwise true whereas $\deg W = \frac{q}{m+1} \deg \Omega^1$ means that $\tau(\rho) = \frac{1}{2m} \int_M \langle f^*\omega_{\mathcal{X}}, \omega \rangle dV = q \text{Vol}(M)$, so that in fact $f^*g_{\mathcal{X}} = qg$ holds everywhere. Proposition 3.2 yields that $f = f_{max}$, up to composition with an isometry of \mathcal{X} .

To get the inequality $\deg W \geq -\frac{q}{m+1} \deg \Omega^1$, consider the map $\gamma : V \rightarrow W \otimes \Omega^1$. We have $\deg V = \deg \text{Ker } \gamma + \deg \text{Im } \gamma$. Since $\text{Ker } \gamma$ is θ -invariant, $\deg \text{Ker } \gamma \leq 0$. By semistability of W , $\deg \text{Im } \gamma \leq \text{rk } \gamma (\frac{1}{q} \deg W + \frac{1}{m} \deg \Omega^1)$. Hence

$$\deg V \leq \frac{q \text{rk } \gamma}{q + \text{rk } \gamma} \frac{\deg \Omega^1}{m} \leq \frac{q}{m+1} \deg \Omega^1$$

with equality if and only if $\text{rk } \gamma = qm$, i.e. γ is generically onto, and $\deg \text{Ker } \gamma = 0$, i.e. $\text{Ker } \gamma$ is a polystable subbundle of E .

Again, the fact that $[\theta, \theta] = 0$ yields that $\beta = 0$, i.e. f is antiholomorphic:

Lemma 3.9. *If $m > 1$ and $\gamma : V \rightarrow W \otimes \Omega^1$ is onto, β vanishes identically.*

Proof of the lemma. Let $X \in T^1$ and $w \in W$. Let $\alpha \in \Omega^1$, $\alpha \neq 0$, be such that $\alpha(X) = 0$. Take $v \in V$ such that $\gamma(v) = w \otimes \alpha$. Then for all $Y \in T^1$, we have on the one hand $\beta(X)\gamma(Y)v = \beta(X)(\alpha(Y)w) = \alpha(Y)\beta(X)w$ and on the other hand $\beta(X)\gamma(Y)v = \beta(Y)\gamma(X)v = \alpha(X)\beta(Y)w = 0$. We may find Y such that $\alpha(Y) \neq 0$, for m is greater than 1. Hence $\beta(X)w = 0$. \square

The rest of the proof goes like in the holomorphic case. \square

3.5. Proof of Proposition 1.2.

We use freely what has been done in Sections 3.2 and 3.3. If the representation ρ is not reductive, we consider its semi-simplification instead. Considering the polystable Higgs bundle $E = V \oplus W$ associated to ρ , we want to prove the inequality

$$|\deg W| \leq \frac{2p}{p+2} \frac{\deg \Omega^1}{m}.$$

The proof again depends on the type of system of Hodge bundles we obtain by deforming E via the \mathbb{C}^* -action.

3.5.1. *System of Hodge bundles of type $V_1 \rightarrow W \rightarrow V_2 \rightarrow 0$.*

As in Paragraph 3.3.1, we need no restriction on the rank of W here. So let $q = \text{rk } W \geq 1$.

We use the method of Viehweg and Zuo [VZ07]. They work with a binary system of Hodge bundles ($V_1 = 0$) so we explain how to adapt their proof to the ternary case. We try to fit to their notations as much as possible. Dualizing the Higgs bundle if necessary, we may suppose that $\deg W > 0$. We also suppose that no subsheaf of V_2 has a slope equal to zero. In fact, each subsheaf of V_2 has non positive slope because $\theta|_{V_2} = 0$ and if its slope is equal to zero, then the Higgs bundle splits as a sum of two polystable Higgs bundles of degree zero with one contained in V_2 .

Let us consider the Harder-Narasimhan filtrations [VZ07, HL97]

$$0 = W^0 \subset W^1 \subset \dots \subset W^{l'} = W$$

and

$$0 = V_2^0 \subset V_2^1 \subset \dots \subset V_2^{l'} = V_2$$

of W and V_2 . Let l be the maximum of all j verifying $\mu(W^j/W^{j-1}) > 0$. Remark that $l \geq 1$ because $\mu(W^1) \geq \mu(W) > 0$.

We construct by induction two sequences

$$0 = j_0 < j_1 < \dots < j_r = l \quad \text{and} \quad 0 = j'_0 < j'_1 < \dots < j'_r \leq l'$$

in the following way:

Suppose that j_{k-1} and j'_{k-1} are defined. If $j_{k-1} < l$, let j'_k be the minimal number with $\beta(W^{j_{k-1}+1}) \subset V_2^{j'_k} \otimes \Omega^1$, and j_k be the maximum of all $j \leq l$ verifying $\beta(W^j) \subset V_2^{j'_k} \otimes \Omega^1$.

Then, we have non trivial morphisms

$$\frac{W^{j_{k-1}+1}}{W^{j_{k-1}}} \longrightarrow \frac{V_2^{j'_k}}{V_2^{j'_{k-1}}} \otimes \Omega^1.$$

Because of the semistability of all involved sheaves, we get

$$\mu\left(\frac{W^{j_{k-1}+1}}{W^{j_{k-1}}}\right) \leq \mu\left(\frac{V_2^{j'_k}}{V_2^{j'_{k-1}}}\right) + \mu(\Omega^1)$$

for each k . We set $E^k = W^{j_k} \oplus V_2^{j'_k}$. The sequence $(E^k)_{0 \leq k \leq r}$ defines a filtration of $W^{j_r} \oplus V_2^{j'_r}$ by Higgs subsheaves, and we denote the successive quotients by $F^k = E^k/E^{k-1} = F_W^k \oplus F_{V_2}^k$, where $F_W^k = W^{j_k}/W^{j_{k-1}}$ and $F_{V_2}^k = V_2^{j'_k}/V_2^{j'_{k-1}}$.

From the properties of the Harder-Narasimhan filtrations, we have

$$\mu(F_W^{k-1}) = \mu\left(\frac{W^{j_{k-1}}}{W^{j_{k-2}}}\right) \geq \mu\left(\frac{W^{j_{k-1}}}{W^{j_{k-1}-1}}\right) > \mu\left(\frac{W^{j_{k-1}+1}}{W^{j_{k-1}}}\right) \geq \mu\left(\frac{W^{j_k}}{W^{j_{k-1}}}\right) = \mu(F_W^k)$$

and

$$\mu(F_{V_2}^k) = \mu\left(\frac{V_2^{j'_k}}{V_2^{j'_{k-1}}}\right) \geq \mu\left(\frac{V_2^{j'_k}}{V_2^{j'_{k-1}-1}}\right) > \mu\left(\frac{V_2^{j'_k+1}}{V_2^{j'_k}}\right) \geq \mu\left(\frac{V_2^{j'_{k+1}}}{V_2^{j'_k}}\right) = \mu(F_{V_2}^{k+1}).$$

In particular, we get for all $1 \leq k \leq r$,

$$\mu(F_W^k) - \mu(F_{V_2}^k) \leq \mu(\Omega^1)$$

and

$$\mu(F_W^1) > \mu(F_W^2) > \dots > \mu(F_W^r) > 0 > \mu(F_{V_2}^1) > \mu(F_{V_2}^2) > \dots > \mu(F_{V_2}^r).$$

Viehweg and Zuo then define the following quantities:

- $c_k = \deg F^k$,
- $\mu_k^W = \mu(F_W^k)$, $\mu_k^{V_2} = \mu(F_{V_2}^k)$,
- $r_k^W = \text{rk}(F_W^k)$,
- $r_k^{V_2} = \text{rk}(F_{V_2}^k) - \frac{c_k}{\mu_k^{V_2}}$.

With these definitions, we can write the above inequalities

$$\mu_k^W - \mu_k^{V_2} \leq \mu(\Omega^1) \quad \text{for all } 1 \leq k \leq r, \quad \text{and} \quad \mu_1^W > \cdots > \mu_r^W > 0 > \mu_1^{V_2} > \cdots > \mu_r^{V_2}.$$

We verify the properties I–IV of Claim 2.2 in [VZ07] (recall that in the present situation, we may have $j_r < l''$):

I. Each $r_k^{V_2} = -\frac{\mu_k^W r_k^W}{\mu_k^{V_2}}$ and hence is positive.

II. Here we need some adaptations. For each $0 \leq k \leq r$, the number $\sum_{i=1}^k c_i$ is non positive because E^k is a Higgs subsheaf. Moreover (if we set $p_1 = \text{rk } V_1$)

$$\begin{aligned} p_1 + \sum_{k=1}^r r_k^{V_2} + \text{rk}(V_2/V_2^{j_r'}) - p &= \sum_{k=1}^r r_k^{V_2} - \sum_{k=1}^r \text{rk}(F_{V_2}^k) = -\sum_{k=1}^r \frac{c_k}{\mu_k^{V_2}} \\ &= -\frac{1}{\mu_r^{V_2}} \left(\sum_{i=1}^r c_i \right) + \sum_{k=1}^{r-1} \frac{\mu_k^{V_2} - \mu_{k+1}^{V_2}}{\mu_k^{V_2} \mu_{k+1}^{V_2}} \left(\sum_{i=1}^k c_i \right) \leq 0. \end{aligned}$$

Thus, $p \geq p_1 + \sum_{k=1}^r r_k^{V_2} + \text{rk}(V_2/V_2^{j_r'}) \geq \sum_{k=1}^r r_k^{V_2}$.

III. By assumption, $\deg V < 0$ and therefore

$$\begin{aligned} \mu(V) &\geq \frac{\sum_{k=1}^r \text{rk}(F_{V_2}^k) \mu_k^{V_2} + \deg(V_2/V_2^{j_r'}) + \deg V_1}{\sum_{k=1}^r r_k^{V_2}} \\ &= \frac{\sum_{k=1}^r r_k^{V_2} \mu_k^{V_2} + \sum_{k=1}^r c_k + \deg(V_2/V_2^{j_r'}) + \deg V_1}{\sum_{k=1}^r r_k^{V_2}} \\ &= \frac{\sum_{k=1}^r r_k^{V_2} \mu_k^{V_2} - \deg(W/W^l)}{\sum_{k=1}^r r_k^{V_2}} \\ &\geq \frac{\sum_{k=1}^r r_k^{V_2} \mu_k^{V_2}}{\sum_{k=1}^r r_k^{V_2}} \end{aligned}$$

(we use $\sum_{k=1}^r c_k + \deg(V_2/V_2^{j_r'}) + \deg V_1 = -\deg(W/W^l) \geq 0$).

IV. From III, we get

$$\mu(W) - \mu(V) \leq \frac{\sum_{k=1}^r r_k^W \mu_k^W}{\sum_{k=1}^r r_k^W} - \frac{\sum_{k=1}^r r_k^{V_2} \mu_k^{V_2}}{\sum_{k=1}^r r_k^{V_2}}.$$

The r.h.s is bounded from above by $\max\{\mu_k^W - \mu_k^{V_2}, 1 \leq k \leq r\}$ (see [VZ07] for the proof of this), hence

$$\frac{p+q}{pq} \deg W = \mu(W) - \mu(V) \leq \frac{\deg \Omega^1}{m}.$$

3.5.2. *System of Hodge bundles of the type $V_1 \rightarrow W_1 \rightarrow V_2 \rightarrow W_2 \rightarrow V_3 \rightarrow 0$.*

If $\beta_1 \neq 0$ and $\gamma_2 \neq 0$, we have exactly as in Section 3.3.2 that

$$|\deg W| \leq \frac{3\text{rk } V_2 + 1}{2(\text{rk } V_2 + 1)} \frac{\deg \Omega^1}{m}$$

using that $\text{rk } \gamma_2 \leq \text{rk } V_2$ instead of $\text{rk } \gamma_2 \leq m$. This inequality is stronger than the one we want to prove here.

If $\beta_1 = 0$ or $\gamma_2 = 0$, the Higgs bundle splits as the sum of two polystable Higgs bundles of degree 0, for example $V_1 \rightarrow W_1 \rightarrow 0$ and $V_2 \rightarrow W_2 \rightarrow V_3 \rightarrow 0$. Since $\text{rk } W_1 = \text{rk } W_2 = 1$, we have by the previous paragraph that

$$|\deg V_1| \leq \frac{\text{rk } V_1}{\text{rk } V_1 + 1} \frac{\deg \Omega^1}{m} \quad \text{and} \quad |\deg (V_2 \oplus V_3)| \leq \frac{\text{rk } V_2 + \text{rk } V_3}{\text{rk } V_2 + \text{rk } V_3 + 1} \frac{\deg \Omega^1}{m}$$

which, in view of the following lemma, gives the result.

Lemma 3.10. *Let p_1, p_2, q_1, q_2 be positive numbers. Let $p = p_1 + p_2$ and $q = q_1 + q_2$. Then*

$$\frac{p_1 q_1}{p_1 + q_1} + \frac{p_2 q_2}{p_2 + q_2} \leq \frac{p q}{p + q}$$

with equality iff $p_1 q_2 = p_2 q_1$.

Proof.

$$\frac{p q}{p + q} - \frac{p_1 q_1}{p_1 + q_1} - \frac{p_2 q_2}{p_2 + q_2} = \frac{(p_1 q_2 - p_2 q_1)^2}{(p + q)(p_1 + q_1)(p_2 + q_2)}.$$

□

4. THE CASE $G = \text{SO}_0(p, 2)$

In this section we prove the main theorem in the case where $G = \text{SO}_0(p, 2)$, the identity component of $\text{SO}(p, 2)$.

4.1. The Hermitian symmetric space $\text{SO}_0(p, 2)/(\text{SO}(p) \times \text{SO}(2))$.

Here, the symmetric space \mathcal{X} is naturally seen as an open subset of the (real) Grassmannian of 2-planes of a real vector space, which makes the complex structure a bit more difficult to understand. Again, details are to be found in [He01, Sat80].

Let $\mathbb{E}_{\mathbb{R}}$ be a real vector space of dimension $p + 2$ ($p \geq 3$), endowed with a non-degenerate quadratic form S of signature $(p, 2)$. The symmetric space \mathcal{X} is defined as the space of all 2-dimensional real subspaces of $\mathbb{E}_{\mathbb{R}}$ on which S is negative-definite. It is an open submanifold of the real Grassmannian of 2-planes of $\mathbb{E}_{\mathbb{R}}$.

Let us fix a 2-plane $\mathbb{W}_{\mathbb{R}}$ of $\mathbb{E}_{\mathbb{R}}$ on which S is negative-definite and let $\mathbb{V}_{\mathbb{R}}$ be its orthogonal complement. We also fix an orientation on $\mathbb{W}_{\mathbb{R}}$.

The group $G = \text{SO}_0(p, 2)$ acts transitively on \mathcal{X} by analytic isomorphisms. The isotropy subgroup K of G at $\mathbb{W}_{\mathbb{R}}$ is identified with the maximal compact subgroup $\text{SO}(p) \times \text{SO}(2)$, and hence \mathcal{X} can be identified with $\text{SO}_0(p, 2)/(\text{SO}(p) \times \text{SO}(2))$.

Let \mathfrak{g} be the Lie algebra of G , $\mathfrak{k} \subset \mathfrak{g}$ the Lie algebra of K and $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ the corresponding Cartan decomposition. Let us fix an orthonormal basis (e_1, \dots, e_p) of $\mathbb{V}_{\mathbb{R}}$ and a direct orthonormal basis (e_{p+1}, e_{p+2}) of $\mathbb{W}_{\mathbb{R}}$ (with respect to $S|_{\mathbb{V}_{\mathbb{R}}}$, respectively $S|_{\mathbb{W}_{\mathbb{R}}}$). We have the following matrix expressions:

$$\mathfrak{k} = \left\{ \left(\begin{array}{cc} X_1 & 0 \\ 0 & X_2 \end{array} \right), X_1 \in M_p(\mathbb{R}), X_2 \in M_2(\mathbb{R}), {}^t X_i = -X_i \ (i = 1, 2) \right\},$$

$$\mathfrak{p} = \left\{ \left(\begin{array}{cc} 0 & A \\ {}^t A & 0 \end{array} \right), A \in M_{p,2}(\mathbb{R}) \right\} \simeq_{\mathbb{R}} \text{Hom}(\mathbb{W}_{\mathbb{R}}, \mathbb{V}_{\mathbb{R}}).$$

The tangent space $T_o \mathcal{X}$ at $o = \mathbb{W}_{\mathbb{R}} \in \mathcal{X}$ will be identified with \mathfrak{p} . The tangent bundle $T\mathcal{X}$ of \mathcal{X} is the bundle $G \times_{\text{Ad}K} \mathfrak{p}$ associated to the K -principal bundle $G \rightarrow \mathcal{X} = G/K$ via the adjoint action of K on \mathfrak{p} .

Since K respects the decomposition $\mathbb{E}_{\mathbb{R}} = \mathbb{V}_{\mathbb{R}} \oplus \mathbb{W}_{\mathbb{R}}$, the vector bundle $E_{\mathbb{R}}$ on \mathcal{X} associated to $G \longrightarrow \mathcal{X}$ via the action of K on $\mathbb{E}_{\mathbb{R}}$ naturally splits as the sum $V_{\mathbb{R}} \oplus W_{\mathbb{R}}$.

Moreover, there exist two natural complex structures belonging to $\mathrm{SO}(S|_{\mathbb{W}_{\mathbb{R}}}) = \mathrm{SO}(2)$ on the 2-dimensional real vector space $\mathbb{W}_{\mathbb{R}}$, and only one that we call I , such that the orientation of the basis (Iw, w) of $\mathbb{W}_{\mathbb{R}}$ coincides with the fixed one (for any non-zero $w \in \mathbb{W}_{\mathbb{R}}$). In the above basis of $\mathbb{W}_{\mathbb{R}}$, the matrix of I is given by

$$I = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

The complex structure I then defines a complex structure on the vector bundle $W_{\mathbb{R}}$ that we still denote by I .

Finally, using the identification $T\mathcal{X} \simeq \mathrm{Hom}(W_{\mathbb{R}}, V_{\mathbb{R}})$, we get the complex structure J on $T\mathcal{X}$: if X is a section of $\mathrm{Hom}(W_{\mathbb{R}}, V_{\mathbb{R}})$, $JX = X \circ I$. The G -invariant Kähler metric $g_{\mathcal{X}}$ on \mathcal{X} is defined at o by

$$g_{\mathcal{X}}(X, Y) = \mathrm{tr}(YX) = 2 \mathrm{tr}({}^tBA), \text{ if } X = \begin{pmatrix} 0 & A \\ {}^tA & 0 \end{pmatrix}, Y = \begin{pmatrix} 0 & B \\ {}^tB & 0 \end{pmatrix} \in \mathfrak{p}.$$

The corresponding Kähler form will still be denoted by $\omega_{\mathcal{X}} = g_{\mathcal{X}}(J., .)$.

Next, we consider the complexifications \mathbb{V} , \mathbb{W} and $\mathbb{E} = \mathbb{V} \oplus \mathbb{W}$ of $\mathbb{V}_{\mathbb{R}}$, $\mathbb{W}_{\mathbb{R}}$ and $\mathbb{E}_{\mathbb{R}}$ respectively. We extend the quadratic form S to \mathbb{E} and the complex structure I to \mathbb{W} by \mathbb{C} -linearity and still denote them by the same letters. Let $\mathbb{W}^{1,0}$ (resp. $\mathbb{W}^{0,1}$) be the eigenspace of I corresponding to the eigenvalue $\sqrt{-1}$ (resp. $-\sqrt{-1}$). These two eigenspaces also are the two isotropic lines in \mathbb{W} for the quadratic form $S|_{\mathbb{W}}$. Moreover, $\mathbb{W}^{0,1}$ may be identified with $(\mathbb{W}^{1,0})^*$ by means of $S|_{\mathbb{W}}$, and \mathbb{V}^* may be identified with \mathbb{V} by means of $S|_{\mathbb{V}}$.

Let us define $e'_{p+1} = \frac{1}{\sqrt{2}}(e_{p+1} + \sqrt{-1}e_{p+2})$ and $e'_{p+2} = \frac{1}{\sqrt{2}}(e_{p+1} - \sqrt{-1}e_{p+2})$. In the sequel, we shall use the basis $(e_1, \dots, e_p, e'_{p+1}, e'_{p+2})$ of \mathbb{E} . The quadratic form S then can be written

$$S = \begin{pmatrix} I_p & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}.$$

In this basis, the complexification $G_{\mathbb{C}}$ of G is the group $\{g \in \mathrm{SL}(p+2, \mathbb{C}) \mid {}^t g S g = S\}$, which is isomorphic to $\mathrm{SO}(p+2, \mathbb{C})$. The complex conjugation of $G_{\mathbb{C}}$ w.r.t. G is given by $\sigma(g) = (I_{p,2} S \bar{g} S I_{p,2})^{-1}$ and the Cartan involution c is the inner automorphism defined by $\alpha = I_{p,2} \in G_{\mathbb{C}}$. The corresponding Cartan decomposition of the Lie algebra of $G_{\mathbb{C}}$ is $\mathfrak{g}_{\mathbb{C}} = \mathfrak{k}_{\mathbb{C}} \oplus \mathfrak{p}_{\mathbb{C}}$ where

$$\mathfrak{p}_{\mathbb{C}} = \left\{ \begin{pmatrix} 0 & C & D \\ {}^tD & 0 & 0 \\ {}^tC & 0 & 0 \end{pmatrix}, C, D \in M_{p,1}(\mathbb{C}) \right\} \simeq_{\mathbb{C}} \mathrm{Hom}(\mathbb{W}_{\mathbb{R}}, \mathbb{V}_{\mathbb{R}}) \otimes \mathbb{C} = \mathrm{Hom}_{\mathbb{C}}(\mathbb{W}, \mathbb{V})$$

and $\mathfrak{k}_{\mathbb{C}}$ is the Lie algebra of $K_{\mathbb{C}}$ which is isomorphic to $\mathrm{SO}(p, \mathbb{C}) \times \mathrm{SO}(2, \mathbb{C})$.

We have

$$\mathfrak{p} = \left\{ \begin{pmatrix} 0 & C & \bar{C} \\ {}^t\bar{C} & 0 & 0 \\ {}^tC & 0 & 0 \end{pmatrix}, C \in M_{p,1}(\mathbb{C}) \right\}$$

and the two eigenspaces of the extended complex structure $J \otimes \mathrm{Id}$ acting on $\mathfrak{p}_{\mathbb{C}}$ are

$$\mathfrak{p}^{1,0} = \left\{ \begin{pmatrix} 0 & C & 0 \\ 0 & 0 & 0 \\ {}^tC & 0 & 0 \end{pmatrix}, C \in M_{p,1}(\mathbb{C}) \right\} \simeq_{\mathbb{C}} \mathrm{Hom}_{\mathbb{C}}(\mathbb{W}^{1,0}, \mathbb{V}) \simeq_{\mathbb{C}} \mathrm{Hom}_{\mathbb{C}}(\mathbb{V}, \mathbb{W}^{0,1})$$

and

$$\mathfrak{p}^{0,1} = \left\{ \begin{pmatrix} 0 & 0 & D \\ {}^t D & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, D \in M_{p,1}(\mathbb{C}) \right\} \simeq_{\mathbb{C}} \text{Hom}_{\mathbb{C}}(\mathbb{W}^{0,1}, \mathbb{V}) \simeq_{\mathbb{C}} \text{Hom}_{\mathbb{C}}(\mathbb{V}, \mathbb{W}^{1,0}).$$

The complexified tangent bundle $T^{\mathbb{C}}\mathcal{X}$ of \mathcal{X} is isomorphic to $G \times_{\text{Ad}K} \mathfrak{p}_{\mathbb{C}} \simeq (G \times_K K_{\mathbb{C}}) \times_{\text{Ad}K_{\mathbb{C}}} \mathfrak{p}_{\mathbb{C}}$, whereas the holomorphic tangent bundle $T^{1,0}\mathcal{X}$ is isomorphic to $G \times_{\text{Ad}K} \mathfrak{p}^{1,0} \simeq (G \times_K K_{\mathbb{C}}) \times_{\text{Ad}K_{\mathbb{C}}} \mathfrak{p}^{1,0}$. There is a natural Hermitian metric on the holomorphic tangent bundle of \mathcal{X} given on $T_o^{1,0}\mathcal{X} \simeq M_{p,1}(\mathbb{C})$ by $h(C, D) = 4\text{tr}({}^t \bar{D}C)$.

The holomorphic sectional curvature for the complex line $\langle X \rangle$ generated by a nonzero $X = \begin{pmatrix} 0 & C & \bar{C} \\ {}^t \bar{C} & 0 & 0 \\ {}^t C & 0 & 0 \end{pmatrix} \in T_o\mathcal{X}$ is given by $K(\langle X \rangle) = -1 + \frac{1}{2} \frac{|{}^t C C|^2}{({}^t \bar{C} C)^2}$. It is clear that $K(\langle X \rangle)$ is pinched between -1 and $-1/2$. The metric $g_{\mathcal{X}}$ is Einstein and its Ricci curvature tensor is $-\frac{p}{2} g_{\mathcal{X}}$.

4.2. Toledo invariant and $\text{SO}_0(p, 2)$ -Higgs bundles.

Let ρ be a representation of a (torsion free) uniform lattice Γ of $\text{SU}(m, 1)$ into $G = \text{SO}_0(p, 2)$, $p \geq 3$. We will assume that ρ is reductive. If it is not, just replace ρ by its semi-simplification ρ_{ss} (see Paragraph 3.3.3) in the following to get the result.

Let $(P_{K_{\mathbb{C}}}, \theta)$ be the G -Higgs principal bundle on M associated to ρ and f as in section 2 and let E be the holomorphic vector bundle on M associated to $P_{K_{\mathbb{C}}}$ via the action of $K_{\mathbb{C}}$ on \mathbb{E} . The bundle E splits holomorphically as the sum of the rank p subbundle $V = P_{K_{\mathbb{C}}} \times_{K_{\mathbb{C}}} \mathbb{V}$ with the rank 2 subbundle $W = P_{K_{\mathbb{C}}} \times_{K_{\mathbb{C}}} \mathbb{W}$. But, in the present situation, we get more structure on W , because $K_{\mathbb{C}}$ also respects the decomposition $\mathbb{W} = \mathbb{W}^{1,0} \oplus \mathbb{W}^{0,1}$. This implies that if we call L the line bundle $P_{K_{\mathbb{C}}} \times_{K_{\mathbb{C}}} \mathbb{W}^{1,0}$, W holomorphically splits as the sum $L \oplus L^{-1}$. Moreover, since $K_{\mathbb{C}}$ preserves $S_{\mathbb{V}}$, we can identify V^* with V . In particular, $\deg V = 0$.

The Higgs field θ is a holomorphic $(1,0)$ -form taking values in the bundle $P_{K_{\mathbb{C}}} \times_{\text{Ad}K_{\mathbb{C}}} \mathfrak{p}_{\mathbb{C}} \simeq \text{Hom}(L, V) \oplus \text{Hom}(L^{-1}, V)$ so that we can write (see also [BGG05])

$$\theta = \begin{pmatrix} 0 & \beta & \gamma \\ {}^t \gamma & 0 & 0 \\ {}^t \beta & 0 & 0 \end{pmatrix}, \text{ where } \begin{cases} \beta : L \longrightarrow V \otimes \Omega^1 \\ \gamma : L^{-1} \longrightarrow V \otimes \Omega^1 \end{cases}$$

Recall that the sheaves Ω^1 and T^1 are semistable (in the usual sense) and the Higgs vector bundle (E, θ) is polystable.

The bundle $f^* T^{1,0}\mathcal{X}$ is isomorphic to the bundle $\text{Hom}(L, V)$ and therefore its degree is given by $-p \deg L$. We obtain that $\tau(\rho) = \frac{4\pi}{m!} \deg L$. Hence the main theorem in this case follows from:

Theorem 4.1. $|\deg L| \leq \frac{1}{m} \deg \Omega^1$, that is $|\tau(\rho)| \leq \frac{m+1}{m} \text{Vol}(M)$. In particular, when $m > 1$, a representation $\rho : \Gamma \longrightarrow \text{SO}_0(p, 2)$ is never maximal.

Proof. We shall denote by θ^2 the morphism of vector bundles

$$\begin{aligned} \theta^2 : T^1 \times T^1 &\longrightarrow \text{End}(E) \\ (X, Y) &\longmapsto \theta(X) \circ \theta(Y) \end{aligned}$$

by ${}^t \gamma \beta$ the morphism

$$\begin{aligned} {}^t \gamma \beta : T^1 \times T^1 &\longrightarrow \text{End}(L) \\ (X, Y) &\longmapsto {}^t \gamma(X) \circ \beta(Y) \end{aligned}$$

etc.

We remark that the system of Hodge bundles obtained after deformation of the Higgs bundle (E, θ) via the \mathbb{C}^* -action on the moduli space is very simple. Indeed, the limiting (E, θ) must verify $\theta^n = 0$ for some n . In particular, as

$$\theta^2 = \begin{pmatrix} \beta^t\gamma + \gamma^t\beta & 0 & 0 \\ 0 & {}^t\gamma\beta & {}^t\gamma\gamma \\ 0 & {}^t\beta\beta & {}^t\beta\gamma \end{pmatrix},$$

the $(2, 2)$ matrix

$$\begin{pmatrix} {}^t\gamma\beta(X, Y) & {}^t\gamma\gamma(X, Y) \\ {}^t\beta\beta(X, Y) & {}^t\beta\gamma(X, Y) \end{pmatrix} = \begin{pmatrix} {}^t\gamma\beta(X, Y) & {}^t\gamma\gamma(X, Y) \\ {}^t\beta\beta(X, Y) & {}^t\beta\gamma(Y, X) \end{pmatrix}$$

(we use $\theta^2(X, Y) = \theta^2(Y, X)$ and in particular ${}^t\beta\gamma(X, Y) = {}^t\beta\gamma(Y, X)$) must be trace free for any $X, Y \in T^1$, which implies ${}^t\gamma\beta = 0$ (and ${}^t\beta\gamma = 0$). Thus

$$\theta^2 = \begin{pmatrix} \beta^t\gamma + \gamma^t\beta & 0 & 0 \\ 0 & 0 & {}^t\gamma\gamma \\ 0 & {}^t\beta\beta & 0 \end{pmatrix}.$$

Suppose now that there exist $X, Y \in T^1$ such that ${}^t\gamma\gamma(X, Y) \neq 0$. For any $X', Y' \in T^1$, ${}^t\beta\beta(X', Y'){}^t\gamma\gamma(X, Y) = 0$ because $\theta^n = 0$, so we conclude that for every $x \in M$, either ${}^t\beta\beta = 0$ or ${}^t\gamma\gamma = 0$ on $T_x^1 \times T_x^1$ and then, by holomorphicity, either ${}^t\beta\beta = 0$ or ${}^t\gamma\gamma = 0$ on $T^1 \times T^1$.

We work on the system of Hodge bundles that we just described. Suppose for example that ${}^t\beta\beta = 0$. Then, the sequence

$$L^{-1} \xrightarrow{\gamma} \text{Im } \gamma \xrightarrow{{}^t\gamma|_{\text{Im } \gamma}} L \xrightarrow{\beta} \text{Im } \beta \longrightarrow 0$$

defines a Higgs subsheaf of (E, θ) .

The bundle $L \otimes T^1$ is semistable and $\text{Im } \beta$ is also a Higgs subsheaf of E , so we have $\mu(L \otimes T^1) \leq \mu(\text{Im } \beta) \leq 0$ and then $\deg L \leq \frac{\deg \Omega^1}{m}$ (if $\beta = 0$, then $\deg L \leq 0$).

Let us consider the maps $\gamma : L^{-1} \otimes T^1 \longrightarrow \text{Im } \gamma$ and ${}^t\gamma|_{\text{Im } \gamma} : \text{Im } \gamma \longrightarrow L \otimes \Omega^1$. We call r and r' their respective rank. By stability, we have

$$\deg \text{Im } \gamma \geq r \left(-\deg L + \frac{1}{m} \deg T^1 \right)$$

and

$$r' \left(\deg L + \frac{1}{m} \deg \Omega^1 \right) \geq \deg \text{Im } {}^t\gamma|_{\text{Im } \gamma} = \deg \text{Im } \gamma - \deg \text{Ker } {}^t\gamma|_{\text{Im } \gamma}.$$

Using the fact that $\deg \text{Ker } {}^t\gamma|_{\text{Im } \gamma} \leq 0$, we immediately get $\deg L \geq -\frac{\deg \Omega^1}{m}$. \square

REFERENCES

- [BGG03] S. B. Bradlow, O. Garcia-Prada and P. B. Gothen, *Surface group representations and $U(p, q)$ -Higgs bundles*, J. Diff. Geom. **64**, 2003, 111-170 2, 3, 5, 13
- [BGG05] S. B. Bradlow, O. Garcia-Prada and P. B. Gothen, *Maximal surface group representations in isometry groups of classical Hermitian symmetric spaces*, Geom. Dedicata **122**, 2006, 185-213 2, 3, 5, 23
- [BI07] M. Burger and A. Iozzi, *Bounded differential forms, generalized Milnor-Wood inequality and an application to deformation rigidity*, Geom. Dedicata **125**, 2007, 1-23 2

- [BIW03] M. Burger, A. Iozzi and A. Wienhard, *Surface group representations with maximal Toledo invariant*, C. R. Acad. Sci. Paris **336**, 2003, 387-390 2
- [BIW06] M. Burger, A. Iozzi and A. Wienhard, *Surface group representations with maximal Toledo invariant*, arXiv:math.DG/0605656 v2, 2006 2
- [CØ01] J.-L. Clerc and B. Ørsted *The Maslov index revisited*, Transform. Groups **6**, 2001, 303-320 2
- [Co88] K. Corlette, *Flat G-bundles with canonical metrics*, J. Diff. Geom. **28**, 1988, 361-382 2, 6
- [DT87] A. Domic and D. Toledo, *The Gromov norm of the Kähler class of symmetric domains*, Math. Ann. **276**, 1987, 425-432 2
- [Eb96] P. Eberlein, *Geometry of nonpositively curved manifolds*, Chicago Lectures in Mathematics, University of Chicago Press, Chicago, IL, 1996 16
- [Go80] W. M. Goldman, *Discontinuous groups and the Euler class*, Thesis, University of California at Berkeley, 1980 2
- [Go88] W. M. Goldman, *Topological components of spaces of representations*, Invent. Math. **93**, no. 3, 1988, 557-607 2
- [GM87] W. M. Goldman and J. J. Millson, *Local rigidity of discrete groups acting on complex hyperbolic space*, Invent. Math. **88**, 1987, 495-520 2
- [He01] S. Helgason, *Differential geometry, Lie groups, and symmetric spaces*, Corrected reprint of the 1978 original, Graduate Studies in Mathematics, 34, American Mathematical Society, Providence, RI, 2001 9, 21
- [Hi87] N. J. Hitchin, *The self-duality equations on a Riemann surface*, Proc London Math. Soc. (3) **55**, 1987, 59-126 2
- [Hi92] N. J. Hitchin, *Lie groups and Teichmüller space*, Topology **31**, 1992, 449-473 2
- [HL97] D. Huybrechts and M. Lehn, *The geometry of moduli spaces of sheaves*, Aspects of Mathematics, E31, Friedr. Vieweg & Sohn, Braunschweig, 1997 14, 19
- [Ko87] S. Kobayashi, *Differential geometry of complex vector bundles*, Princeton University Press, 1987 17
- [KM08] V. Koziarz and J. Maubon, *Harmonic maps and representations of non-uniform lattices of $PU(m, 1)$* , Ann. Inst. Fourier (Grenoble) **58**, no. 2, 2008, 507-558 2
- [La91] F. Labourie, *Existence d'applications harmoniques tordues à valeurs dans les variétés à courbure négative*, Proc. Amer. Math. Soc. **111**, no. 3, 1991, 877-882 16
- [Ro80] H. L. Royden, *The Ahlfors-Schwarz lemma in several complex variables*, Comment. Math. Helvetici **55**, 1980, 547-558 11
- [Sam86] J. H. Sampson, *Applications of harmonic maps to Kähler geometry*, Contemp. Math. **49**, 1986, 125-133 7
- [Sat80] I. Satake, *Algebraic structures of symmetric domains*, Kanô Memorial Lectures, 4, Iwanami Shoten, Tokyo; Princeton University Press, Princeton, N.J., 1980 9, 11, 21
- [Si88] C. T. Simpson, *Constructing variations of Hodge structure using Yang-Mills theory and applications to uniformization*, J. Amer. Math. Soc. **1**, 1988, 867-918 2, 5, 7, 8
- [Si92] C. T. Simpson, *Higgs bundles and local systems*, Inst. Hautes Études Sci. Publ. Math. **75**, 1992, 5-95 2, 5, 7, 8, 9
- [Si94a] C. T. Simpson, *Moduli of representations of the fundamental group of a smooth projective variety I*, Inst. Hautes Études Sci. Publ. Math. **79**, 1994, 47-129 2, 5, 8
- [Si94b] C. T. Simpson, *Moduli of representations of the fundamental group of a smooth projective variety II*, Inst. Hautes Études Sci. Publ. Math. **80**, 1994, 5-79 2, 5, 8
- [To79] D. Toledo, *Harmonic mappings of surfaces to certain Kähler manifolds*, Math. Scand. **45**, 1979, 13-26 2
- [To89] D. Toledo, *Representations of surface groups in complex hyperbolic space*, J. Diff. Geom. **29**, 1989, 125-133 2
- [VZ07] E. Viehweg and K. Zuo, *Arakelov inequalities and the uniformization of certain rigid Shimura varieties*, J. Differential Geom. **77**, no. 2, 2007, 291-352 4, 14, 19, 20
- [Xi00] E. Z. Xia, *The moduli of flat $PU(2, 1)$ structures over Riemann surfaces*, Pacific J. Math. **193**, 2000, 231-256 3, 13

INSTITUT ELIE CARTAN, UNIVERSITÉ HENRI POINCARÉ, B. P. 239, F-54506 VANDŒUVRE-LÈS-NANCY CEDEX, FRANCE

E-mail address: koziarz@iecn.u-nancy.fr

E-mail address: maubon@iecn.u-nancy.fr