

# A GENERIC PROPERTY OF FAMILIES OF LAGRANGIAN SYSTEMS

PATRICK BERNARD AND GONZALO CONTRERAS

ABSTRACT. We prove that a generic lagrangian has finitely many minimizing measures for every cohomology class.

Annals of Maths, to appear.

## 1. INTRODUCTION

Let  $M$  be a compact boundaryless smooth manifold.

Let  $\mathbb{T}$  be either the group  $(\mathbb{R}/\mathbb{Z}, +)$  or the trivial group  $(\{0\}, +)$ .

A *Tonelli Lagrangian* is a  $C^2$  function  $L : \mathbb{T} \times TM \rightarrow \mathbb{R}$  such that

- The restriction to each fiber of  $\mathbb{T} \times TM \rightarrow \mathbb{T} \times M$  is a *convex* function.
- It is fiberwise *superlinear*:

$$\lim_{|\theta| \rightarrow +\infty} L(t, \theta)/|\theta| = +\infty, \quad (t, \theta) \in \mathbb{T} \times TM.$$

- The Euler-Lagrange equation

$$\frac{d}{dt} L_v = L_x$$

defines a *complete* flow  $\varphi : \mathbb{R} \times (\mathbb{T} \times TM) \rightarrow \mathbb{T} \times TM$ .

We say that a Tonelli Lagrangian  $L$  is *strong Tonelli* if  $L + u$  is a Tonelli Lagrangian for each  $u \in C^\infty(\mathbb{T} \times M, \mathbb{R})$ . When  $\mathbb{T} = \{0\}$  we say that the lagrangian is *autonomous*.

Let  $\mathcal{P}(L)$  be the set of Borel probability measures on  $\mathbb{T} \times TM$  which are invariant under the Euler-Lagrange flow  $\varphi$ . The action functional  $A_L : \mathcal{P}(L) \rightarrow \mathbb{R} \cup \{+\infty\}$  is defined as

$$A_L(\mu) := \langle L, \mu \rangle := \int_{\mathbb{T} \times TM} L \, d\mu.$$

The functional  $A_L$  is lower semi-continuous and the minimizers of  $A_L$  on  $\mathcal{P}(L)$  are called *minimizing measures*. The ergodic components of a minimizing measure are also minimizing, and they are mutually singular, so that the set  $\mathfrak{M}(L)$  of minimizing measures is a simplex whose extremal points are the ergodic minimizing measures.

In general, the simplex  $\mathfrak{M}(L)$  may be of infinite dimension. The goal of the present paper is to prove that this is a very exceptional phenomenon. The first results in that direction were obtained by Mañé in [4]. His paper has been very influential to our work.

---

Gonzalo Contreras was partially supported by CONACYT-México grant # 46467-F.

We say that a property is *generic* in the sense of Mañé if, for each strong Tonelli Lagrangian  $L$ , there exists a residual subset  $\mathcal{O} \subset C^\infty(\mathbb{T} \times M, \mathbb{R})$  such that the property holds for all the Lagrangians  $L - u, u \in \mathcal{O}$ . A set is called residual if it is a countable intersection of open and dense sets. We recall which topology is used on  $C^\infty(\mathbb{T} \times M, \mathbb{R})$ . Denoting by  $\|u\|_k$  the  $C^k$ -norm of a function  $u : \mathbb{T} \times M \rightarrow \mathbb{R}$ , define

$$\|u\|_\infty := \sum_{k \in \mathbb{N}} \frac{\arctan(\|u\|_k)}{2^k}.$$

Note that  $\|\cdot\|_\infty$  is not a norm. Endow the space  $C^\infty(\mathbb{T} \times M, \mathbb{R})$  with the translation-invariant metric  $\|u - v\|_\infty$ . This metric is complete, hence the Baire property holds: any residual subset of  $C^\infty(\mathbb{T} \times M, \mathbb{R})$  is dense.

**Theorem 1.** *Let  $A$  be a finite dimensional convex family of strong Tonelli Lagrangians. Then there exists a residual subset  $\mathcal{O}$  of  $C^\infty(\mathbb{T} \times M, \mathbb{R})$  such that,*

$$u \in \mathcal{O}, \quad L \in A \quad \implies \quad \dim \mathfrak{M}(L - u) \leq \dim A.$$

*In other words, there exist at most  $1 + \dim A$  ergodic minimizing measures of  $L - u$ .*

The main result of Mañé in [4] is that having a unique minimizing measure is a generic property. This corresponds to the case where  $A$  is a point in our statement. Our generalization of Mañé's result is motivated by the following construction due to John Mather:

We can view a 1-form on  $M$  as a function on  $TM$  which is linear on the fibers. If  $\lambda$  is closed, the Euler-Lagrange equation of the Lagrangian  $L - \lambda$  is the same as that of  $L$ . However, the minimizing measures of  $L - \lambda$ , are not the same as the minimizing measures of  $L$ . Mather proves in [5] that the set  $\mathfrak{M}(L - \lambda)$  of minimizing measures of the lagrangian  $L - \lambda$  depends only on the cohomology class  $c$  of  $\lambda$ . If  $c \in H^1(M, \mathbb{R})$  we write  $\mathfrak{M}(L - c) := \mathfrak{M}(L - \lambda)$ , where  $\lambda$  is a closed form of cohomology  $c$ .

It turns out that important applications of Mather theory, such as the existence of orbits wandering in phase space, require understanding not only of the set  $\mathfrak{M}(L)$  of minimizing measures for a fixed or generic cohomology classes but of the set of all Mather minimizing measures for every  $c \in H^1(M, \mathbb{R})$ . The following corollaries are crucial for these applications.

**Corollary 2.** *The following property is generic in the sense of Mañé:*

*For all  $c \in H^1(M, \mathbb{R})$ , there are at most  $1 + \dim H^1(M, \mathbb{R})$  ergodic minimizing measures of  $L - c$ .*

We say that a property is of infinite codimension if, for each finite dimensional convex family  $A$  of strong Tonelli Lagrangians, there exists a residual subset  $\mathcal{O}$  in  $C^\infty(\mathbb{T} \times M, \mathbb{R})$  such that none of the Lagrangians  $L - u, L \in A, u \in \mathcal{O}$  satisfy the property.

**Corollary 3.** *The following property is of infinite codimension:*

*There exists  $c \in H^1(M, \mathbb{R})$ , such that  $L - c$  has infinitely many ergodic minimizing measures.*

Another important issue concerning variational methods for Arnold diffusion questions is the total disconnectedness of the quotient Aubry set. John Mather proves in [7, § 3] that the quotient Aubry set  $\overline{\mathcal{A}}$  of any Tonelli lagrangian on  $\mathbb{T} \times TM$  with  $\mathbb{T} = \mathbb{R}/\mathbb{Z}$  and  $\dim M \leq 2$  (or with  $\mathbb{T} = \{0\}$  and  $\dim M \leq 3$ ) is totally disconnected. See [7] for its definition.

The elements of the quotient Aubry set are called *static classes*. They are disjoint subsets of  $\mathbb{T} \times TM$  and each static class supports at least one ergodic minimizing measure. We then get

**Corollary 4.** *The following property is generic in the sense of Mañé:*

*For all  $c \in H^1(M, \mathbb{R})$  the quotient Aubry set  $\overline{\mathcal{A}}_c$  of  $L - c$  has at most  $1 + \dim H^1(M, \mathbb{R})$  elements.*

## 2. ABSTRACT RESULTS

Assume that we are given

- Three topological vector spaces  $E, F, G$ .
- A continuous linear map  $\pi : F \rightarrow G$ .
- A bilinear pairing  $\langle u, \nu \rangle : E \times G \rightarrow \mathbb{R}$ .
- Two metrizable convex compact subsets  $H \subset F$  and  $K \subset G$  such that  $\pi(H) \subset K$ .

Suppose that

- (i) The map

$$E \times K \ni (u, \nu) \longmapsto \langle u, \nu \rangle$$

is continuous.

We will also denote  $\langle u, \pi(\mu) \rangle$  by  $\langle u, \mu \rangle$  when  $\mu \in H$ . Observe that each element  $u \in E$  gives rise to a linear functional on  $F$

$$F \ni \mu \longmapsto \langle u, \mu \rangle$$

which is continuous on  $H$ . We shall denote by  $H^*$  the set of affine and continuous functions on  $H$  and use the same symbol  $u$  for an element of  $E$  and for the element  $u \longmapsto \langle u, \mu \rangle$  of  $H^*$  which is associated to it.

- (ii) The compact  $K$  is separated by  $E$ . This means that, if  $\eta$  and  $\nu$  are two different points of  $K$ , then there exists a point  $u$  in  $E$  such that  $\langle u, \eta - \nu \rangle \neq 0$ .

Note that the topology on  $K$  is then the weak topology associated to  $E$ . A sequence  $\eta_n$  of elements of  $K$  converges to  $\eta$  if and only if we have  $\langle u, \eta_n \rangle \longrightarrow \langle u, \eta \rangle$  for each  $u \in E$ . We shall, for notational conveniences, fix once and for all a metric  $d$  on  $K$ .

- (iii)  $E$  is a Frechet space. It means that  $E$  is a topological vector space whose topology is defined by a translation-invariant metric, and that  $E$  is complete for this metric.

Note then that  $E$  has the Baire property. We say that a subset is residual if it is a countable intersection of open and dense sets. The Baire property says that any residual subset of  $E$  is dense.

Given  $L \in H^*$  denote by

$$M_H(L) := \arg \min_H L$$

the set of points  $\mu \in H$  which minimize  $L|_H$ , and by  $M_K(L)$  the image  $\pi(M_H(L))$ . These are compact convex subsets of  $H$  and  $K$ .

Our main abstract result is:

**Theorem 5.** *For every finite dimensional affine subspace  $A$  of  $H^*$ , there exists a residual subset  $\mathcal{O}(A) \subset E$  such that, for all  $u \in \mathcal{O}(A)$  and all  $L \in A$ , we have*

$$(1) \quad \dim M_K(L - u) \leq \dim A.$$

**Proof:** We define the  $\varepsilon$ -neighborhood  $V_\varepsilon$  of a subset  $V$  of  $K$  as the union of all the open balls in  $K$  which have radius  $\varepsilon$  and are centered in  $V$ . Given a subset  $D \subset A$ , a positive number  $\varepsilon$ , and a positive integer  $k$ , denote by  $\mathcal{O}(D, \varepsilon, k) \subset E$  the set of points  $u \in E$  such that, for each  $L \in D$ , the convex set  $M_K(L - u)$  is contained in the  $\varepsilon$ -neighborhood of some  $k$ -dimensional convex subset of  $K$ .

We shall prove that the theorem holds with

$$\mathcal{O}(A) = \bigcap_{\varepsilon > 0} \mathcal{O}(A, \varepsilon, \dim A).$$

If  $u$  belongs to  $\mathcal{O}(A)$ , then (1) holds for every  $L \in A$ . Otherwise, for some  $L \in A$ , the convex set  $M_K(L - u)$  would contain a ball of dimension  $\dim A + 1$ , and, if  $\varepsilon$  is small enough, such a ball is not contained in the  $\varepsilon$ -neighborhood of any convex set of dimension  $\dim A$ .

So we have to prove that  $\mathcal{O}(A)$  is residual. In view of the Baire property, it is enough to check that, for any compact subset  $D \subset A$  and any positive  $\varepsilon$ , the set  $\mathcal{O}(D, \varepsilon, \dim A)$  is open and dense. We shall prove in 2.1 that it is open, and in 2.2 that it is dense.

□

### 2.1. Open.

We prove that, for any  $k \in \mathbb{Z}^+$ ,  $\varepsilon > 0$  and any compact  $D \subset A$ , the set  $\mathcal{O}(D, \varepsilon, k) \subset E$  is open. We need a Lemma.

**Lemma 6.** *The set-valued map  $(L, u) \mapsto M_H(L - u)$  is upper semi-continuous on  $A \times E$ . This means that for any open subset  $U$  of  $H$ , the set*

$$\{(L, u) \in A \times E : M_H(L - u) \subset U\} \subset A \times E$$

*is open in  $A \times E$ . Consequently, the set-valued map  $(L, u) \mapsto M_K(L - u)$  is also upper semi-continuous.*

**Proof:** This is a standard consequence of the continuity of the map

$$A \times E \times H \ni (L, u, \mu) \longmapsto (L - u)(\mu) = L(\mu) - \langle u, \mu \rangle.$$

□

Now let  $u_0$  be a point of  $\mathcal{O}(D, \varepsilon, k)$ . For each  $L \in D$ , there exists a  $k$ -dimensional convex set  $V \subset K$  such that  $M_K(L - u_0) \subset V_\varepsilon$ . In other words, the open sets of the form

$$\{(L, u) \in D \times E : M_H(L - u) \subset V_\varepsilon\} \subset D \times E,$$

where  $V$  is some  $k$ -dimensional convex subset of  $K$ , cover the compact set  $D \times \{u_0\}$ . So there exists a finite subcovering of  $D \times \{u_0\}$  by open sets of the form  $\Omega_i \times U_i$ , where  $\Omega_i$  is an open set in  $A$  and  $U_i \subset \mathcal{O}(\Omega_i, \varepsilon, k)$  is an open set in  $E$  containing  $u_0$ . We conclude that the open set  $\cap U_i$  is contained in  $\mathcal{O}(D, \varepsilon, k)$ , and contains  $u_0$ . This ends the proof.

□

## 2.2. Dense.

We prove the density of  $\mathcal{O}(A, \varepsilon, \dim A)$  in  $E$  for  $\varepsilon > 0$ . Let  $w$  be a point in  $E$ . We want to prove that  $w$  is in the closure of  $\mathcal{O}(A, \varepsilon, \dim A)$ .

**Lemma 7.** *There exists an integer  $m$  and a continuous map*

$$T_m = (w_1, \dots, w_m) : K \longrightarrow \mathbb{R}^m,$$

with  $w_i \in E$  such that

$$(2) \quad \forall x \in \mathbb{R}^m \quad \text{diam } T_m^{-1}(x) < \varepsilon,$$

where the diameter is taken for the distance  $d$  on  $K$ .

**Proof:** In  $K \times K$ , to each element  $w \in E$  we associate the open set

$$U_w = \{(\eta, \mu) \in K \times K : \langle w, \eta - \mu \rangle \neq 0\}.$$

Since  $E$  separates  $K$ , the open sets  $U_w, w \in E$  cover the complement of the diagonal in  $K \times K$ . Since this complement is open in the separable metrizable set  $K \times K$ , we can extract a countable subcovering from this covering. So we have a sequence  $U_{w_k}$ , with  $w_k \in E$ , which covers the complement of the diagonal in  $K \times K$ . This amounts to say that the sequence  $w_k$  separates  $K$ . Defining  $T_m = (w_1, \dots, w_m)$ , we have to prove that (2) holds for  $m$  large enough. Otherwise, we would have two sequences  $\eta_m$  and  $\mu_m$  in  $K$  such that

$$T_m(\mu_m) = T_m(\eta_m) \quad \text{and} \quad d(\mu_m, \eta_m) \geq \varepsilon.$$

By extracting a subsequence, we can assume that the sequences  $\mu_m$  and  $\eta_m$  have different limits  $\mu$  and  $\eta$ , which satisfy  $d(\eta, \mu) \geq \varepsilon$ . Take  $m$  large enough, so that  $T_m(\eta) \neq T_m(\mu)$ . Such a value of  $m$  exists because the linear forms  $w_k$  separate  $K$ . We have that

$$T_m(\mu_k) = T_m(\eta_k) \quad \text{for} \quad k \geq m.$$

Hence at the limit  $T_m(\eta) = T_m(\mu)$ . This is a contradiction.

□

Define the function  $F_m : A \times \mathbb{R}^m \rightarrow \mathbb{R} \cup \{+\infty\}$  as

$$F_m(L, x) := \min_{\substack{\mu \in H \\ T_m \circ \pi(\mu) = x}} (L - w)(\mu),$$

when  $x \in T_m(\pi(H))$  and  $F_m(L, x) = +\infty$  if  $x \in \mathbb{R}^m \setminus T_m(\pi(H))$ . For  $y = (y_1, \dots, y_m) \in \mathbb{R}^m$ , let

$$M_m(L, y) := \arg \min_{x \in \mathbb{R}^m} [F_m(L, x) - y \cdot x] \subset \mathbb{R}^m$$

be the set of points which minimize the function  $x \mapsto F_m(L, x) - y \cdot x$ . We have that

$$M_K(L - w - y_1 w_1 - \dots - y_m w_m) \subset T_m^{-1}(M_m(L, y)).$$

Let

$$\mathcal{O}_m(A, \dim A) := \{ y \in \mathbb{R}^m \mid \forall L \in A : \dim M_m(L, y) \leq \dim A \}.$$

From Lemma 7 it follows that

$$y \in \mathcal{O}_m(A, \dim A) \implies w + y_1 w_1 + \dots + y_m w_m \in \mathcal{O}(A, \varepsilon, \dim A).$$

Therefore, in order to prove that  $w$  is in the closure of  $\mathcal{O}(A, \varepsilon, \dim A)$ , it is enough to prove that  $0$  is in the closure of  $\mathcal{O}_m(A, \dim A)$ , which follows from the next proposition.

**Proposition 8.** *The set  $\mathcal{O}_m(A, \dim A)$  is dense in  $\mathbb{R}^m$ .*

**Proof:** Consider the Legendre transform of  $F_m$  with respect to the second variable,

$$\begin{aligned} G_m(L, y) &= \max_{x \in \mathbb{R}^m} y \cdot x - F_m(L, x) \\ &= \max_{\mu \in H} (w + y_1 w_1 + \dots + y_m w_m, \mu) - L(\mu). \end{aligned}$$

It follows from this second expression that the function  $G_m$  is convex and finite-valued, hence continuous on  $A \times \mathbb{R}^m$ .

Consider the set  $\tilde{\Sigma} \subset A \times \mathbb{R}^m$  of points  $(L, y)$  such that  $\dim \partial G_m(L, y) \geq \dim A + 1$ , where  $\partial G_m$  is the subdifferential of  $G_m$ . It is known, see the appendix, that this set has Hausdorff dimension at most

$$(m + \dim A) - (\dim A + 1) = m - 1.$$

Consequently, the projection  $\Sigma$  of the set  $\tilde{\Sigma}$  on the second factor  $\mathbb{R}^m$  also has Hausdorff dimension at most  $m - 1$ . Therefore, the complement of  $\Sigma$  is dense in  $\mathbb{R}^m$ . So it is enough to prove that

$$y \notin \Sigma \implies \forall L \in A : \dim M_m(L, y) \leq \dim A.$$

Since we know by definition of  $\Sigma$  that  $\dim \partial G_m(L, y) \leq \dim A$ , it is enough to observe that

$$\dim M_m(L, y) \leq \dim \partial G_m(L, y).$$

The last inequality follows from the fact that the set  $M_m(L, y)$  is the subdifferential of the convex function

$$\mathbb{R}^m \ni z \mapsto G_m(L, z)$$

at the point  $y$ .

### 3. APPLICATION TO LAGRANGIAN DYNAMICS

Let  $C$  be the set of continuous functions  $f : \mathbb{T} \times TM \rightarrow \mathbb{R}$  with linear growth, i.e.

$$\|f\|_\ell := \sup_{(t,\theta) \in \mathbb{T} \times TM} \frac{|f(t,\theta)|}{1+|\theta|} < +\infty,$$

endowed with the norm  $\|\cdot\|_\ell$ .

We apply Theorem 5 to the following setting:

- $F = C^*$  is the vector space of continuous linear functionals  $\mu : C \rightarrow \mathbb{R}$  provided with the weak- $\star$  topology. Recall that

$$\lim_n \mu_n = \mu \iff \lim_n \mu_n(f) = \mu(f), \quad \forall f \in C.$$

- $E = C^\infty(\mathbb{T} \times M, \mathbb{R})$  provided with the  $C^\infty$  topology.
- $G$  is the vector space of finite Borel signed measures on  $TM$ , or equivalently the set of continuous linear forms on  $C^0(\mathbb{T} \times M, \mathbb{R})$ , provided with the weak- $\star$  topology.
- The pairing  $E \times G \rightarrow \mathbb{R}$  is given by integration:

$$\langle u, \nu \rangle = \int_{\mathbb{T} \times M} u \, d\nu.$$

- The continuous linear map  $\pi : F \rightarrow G$  is induced by the projection  $\mathbb{T} \times TM \rightarrow \mathbb{T} \times M$ .
- The compact  $K \subset G$  is the set of Borel probability measures on  $\mathbb{T} \times M$ , provided with the weak- $\star$  topology. Observe that  $K$  is separated by  $E$ .
- The compact  $H_n \subset F$  is the set of holonomic probability measures which are supported on

$$B_n := \{(t, \theta) \in \mathbb{T} \times TM \mid |\theta| \leq n\}.$$

Holonomic probabilities are defined as follows: Given a  $C^1$  curve  $\gamma : \mathbb{R} \rightarrow M$  of period  $T \in \mathbb{N}$  define the element  $\mu_\gamma$  of  $F$  by

$$\langle f, \mu_\gamma \rangle = \frac{1}{T} \int_0^T f(s, \gamma(s), \dot{\gamma}(s)) \, ds$$

for each  $f \in C$ . Let

$$\Gamma := \{ \mu_\gamma \mid \gamma \in C^1(\mathbb{R}, M) \text{ is periodic of integral period} \} \subset F.$$

The set  $\mathcal{H}$  of holonomic probabilities is the closure of  $\Gamma$  in  $F$ . One can see that  $\mathcal{H}$  is convex (cf. Mañé [4, prop. 1.1(a)]). The elements  $\mu$  of  $\mathcal{H}$  satisfy  $\langle 1, \mu \rangle = 1$  therefore we have  $\pi(\mathcal{H}) \subset K$ .

Note that each Tonelli Lagrangian  $L$  gives rise to an element of  $H_n^*$ .

Let  $\mathfrak{M}(L)$  be the set of minimizing measures for  $L$  and let  $\text{supp } \mathfrak{M}(L)$  be the union of their supports. Recalling that we have defined  $M_{H_n}(L)$  as the set of measures  $\mu \in H_n$  which minimize the action  $\int L \, d\mu$  on  $H_n$ , we have:

**Lemma 9.** *If  $L$  is a Tonelli lagrangian then there exists  $n \in \mathbb{N}$  such that*

$$\dim \pi(M_{H_n}(L)) = \dim \mathfrak{M}(L).$$

**Proof:**

Birkhoff theorem implies that  $\mathfrak{M}(L) \subset \mathcal{H}$  (cf. Mañé [4, prop. 1.1.(b)]). In [5, Prop. 4, p. 185] Mather proves that  $\text{supp } \mathfrak{M}(L)$  is compact, therefore  $\mathfrak{M}(L) \subset H_n$  for some  $n \in \mathbb{N}$ .

In [4, §1] Mañé proves that minimizing measures are also all the minimizers of the action functional  $A_L(\mu) = \int L d\mu$  on the set of holonomic measures, therefore  $\mathfrak{M}(L) = M_{H_n}(L)$  for some  $n \in \mathbb{N}$ .

In [5, Th. 2, p. 186] Mather proves that the restriction  $\text{supp } \mathfrak{M}(L) \rightarrow M$  of the projection  $TM \rightarrow M$  is injective. Therefore the linear map  $\pi : \mathfrak{M}(L) \rightarrow G$  is injective, so that  $\dim \pi(M_{H_n}(L)) = \dim \pi(\mathfrak{M}(L)) = \dim \mathfrak{M}(L)$ .  $\square$

**Proof of Theorem 1.**

Given  $n \in \mathbb{N}$  apply Theorem 5 and obtain a residual subset  $\mathcal{O}_n(A) \subset E$  such that

$$L \in A, \quad u \in \mathcal{O}_n(A) \implies \dim \pi(M_{H_n}(L - u)) \leq \dim A.$$

Let  $\mathcal{O}(A) = \bigcap_n \mathcal{O}_n(A)$ . By the Baire property  $\mathcal{O}(A)$  is residual. We have that

$$L \in A, \quad u \in \mathcal{O}(A), \quad n \in \mathbb{N} \implies \dim \pi(M_{H_n}(L - u)) \leq \dim A.$$

Then by Lemma 9,  $\dim \mathfrak{M}(L - u) \leq \dim A$  for all  $L \in A$  and all  $u \in \mathcal{O}(A)$ .  $\square$

## APPENDIX A. CONVEX FUNCTIONS

Given a convex function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  and  $x \in \mathbb{R}^n$ , define its subdifferential as

$$\partial f(x) := \{ \ell : \mathbb{R}^n \rightarrow \mathbb{R} \text{ linear} \mid f(y) \geq f(x) + \ell(y - x), \forall y \in \mathbb{R}^n \}.$$

Then the sets  $\partial f(x) \subset \mathbb{R}^n$  are convex. If  $k \in \mathbb{N}$ , let

$$\Sigma_k(f) := \{ x \in \mathbb{R}^n \mid \dim \partial f(x) \geq k \}.$$

The following result is standard.

**Proposition 10.** *If  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is a convex function then for all  $0 \leq k \leq n$  the Hausdorff dimension  $HD(\Sigma_k(f)) \leq n - k$ .*

We recall here an elegant proof due to Ambrosio and Alberti, see [1]. Note that much more can be said on the structure of  $\Sigma_k$ , see [2, 9] for example.

By adding  $|x|^2$  if necessary (which does not change  $\Sigma_k$ ) we can assume that  $f$  is superlinear and that

$$(3) \quad f(y) \geq f(x) + \ell(y - x) + \frac{1}{2} |y - x|^2 \quad \forall x, y \in \mathbb{R}^n, \quad \forall \ell \in \partial f(x).$$

**Lemma 11.**  $\ell \in \partial f(x), \quad \ell' \in \partial f(x') \implies |x - x'| \leq \|\ell - \ell'\|.$



**Proof:** From inequality (3) we have that

$$\begin{aligned} f(x') &\geq f(x) + \ell(x' - x) + \frac{1}{2} |x' - x|^2, \\ f(x) &\geq f(x') + \ell'(x - x') + \frac{1}{2} |x - x'|^2. \end{aligned}$$

Then

$$\begin{aligned} (4) \quad &0 \geq (\ell' - \ell)(x - x') + |x - x'|^2 \\ (5) \quad &\|\ell - \ell'\| |x - x'| \geq (\ell - \ell')(x - x') \geq |x - x'|^2. \end{aligned}$$

Therefore  $\|\ell - \ell'\| \geq |x - x'|$ . □

Since  $f$  is superlinear, the subdifferential  $\partial f$  is surjective and we have:

**Corollary 12.** *There exists a Lipschitz function  $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$  such that*

$$\ell \in \partial f(x) \implies x = F(\ell).$$

**Proof of Proposition 10:**

Let  $A_k$  be a set with  $HD(A_k) = n - k$  such that  $A_k$  intersects any convex subset of dimension  $k$ . For example

$$A_k = \{x \in \mathbb{R}^n \mid x \text{ has at least } k \text{ rational coordinates}\}.$$

Observe that

$$x \in \Sigma_k \implies \partial f(x) \text{ intersects } A_k \implies x \in F(A_k).$$

Therefore  $\Sigma_k \subset F(A_k)$ . Since  $F$  is Lipschitz, we have that  $HD(\Sigma_k) \leq HD(A_k) = n - k$ . □

## REFERENCES

1. Giovanni Alberti; Luigi Ambrosio, *A geometrical approach to monotone functions in  $R^n$* . Math. Z. 230 (1999), no. 2, 259–316.
2. Giovanni Alberti, *On the structure of singular sets of convex functions*, Calc. Var. & Partial Differential Equations **2** (1994), no. 1, 17–27.
3. Ricardo Mañé, *On the minimizing measures of Lagrangian dynamical systems*, Nonlinearity **5** (1992), no. 3, 623–638.
4. Ricardo Mañé, *Generic properties and problems of minimizing measures of Lagrangian systems*, Nonlinearity **9** (1996), no. 2, 273–310.
5. John N. Mather, *Action minimizing invariant measures for positive definite Lagrangian systems*, Math. Z. **207** (1991), no. 2, 169–207.
6. John N. Mather, *Variational construction of connecting orbits*, Ann. Inst. Fourier (Grenoble) **43** (1993), no. 5, 1349–1386.
7. John N. Mather, *A property of compact, connected, laminated subsets of manifolds*, Ergodic Theory Dynam. Systems **22** (2002), no. 5, 1507–1520.
8. John N. Mather, *Examples of Aubry sets*. Ergodic Theory Dyn. Syst. **24** (2004), no. 5, 1667–1723.
9. Luděk Zajček, *On the points of multiplicity of monotone operators*. Comment. Math. Univ. Carolinae **19** (1978), no. 1, 179–189.

CEREMADE, UNIVERSITÉ DE PARIS DAUPHINE, PL. DU MARÉCHAL DE LATTRE DE TASSIGNY, 75775  
PARIS CEDEX 16, FRANCE,

*E-mail address:* `patrick.bernard@ceremade.dauphine.fr`

CIMAT, P.O. Box 402, 36.000 GUANAJUATO, GTO, MÉXICO.

*E-mail address:* `gonzalo@ciamat.mx`